arXiv:1211.0936v1 [math.RA] 5 Nov 2012

# THE HOMOGENIZED ENVELOPING ALGEBRA OF THE LIE ALGEBRA $s\ell(2,\mathbb{C})$.

ROBERTO MARTINEZ-VILLA

Abstract. In this paper we study the homogenized algebra $B$ of the enveloping algebra $U$ of the Lie algebra s$\ell(2,\mathbb{C})$. We look first to connections between the category of graded left $B$- modules and the category of $U$-modules, then we prove $B$ is Koszul and Artin-Schelter regular of global dimension four, hence its Yoneda algebra $B^!$ is selfinjective of radical five zero, the structure of $B^!$ is given. We describe next the category of homogenized Verma modules, which correspond to the lifting to $B$ of the usual Verma modules over $U$, and prove that such modules are Koszul of projective dimension two. It was proved in [MZ] that all graded stable components of a selfinjective Koszul algebra are of type $ZA_\infty$, we characterize here the graded $B^!$-modules corresponding under Koszul duality to the homogenized Verma modules, and prove that they are located at the mouth of a regular component, in this way we obtain a family of components over a wild algebra indexed by $\mathbb{C}$.

The paper ends with the description of a family of weight modules over $B$ which corresponds to the weight modules over $U$ and with a description of the category of $B$ - modules corresponding to the Gelfand's category $\mathcal{O}$ of $U$-modules .

## 1. Introduction

In the papers $[MMo1]$,[MMo2] we considered the homogenized Weyl algebra $B_n$ of the Weyl algebra $A_n$ [Co]. We proved that such algebra is Artin-Schelter regular and Koszul and explored the relation between the category of graded $B_n$-modules and the category of $A_n$-modules. Since the connection between these two categories is well understood, we can study the finitely generated graded $B_n$- modules instead of the finitely generated $A_n$-modules. In those papers we applied Koszul duality and related the finitely graded $B_n$- modules with the finitely generated graded modules over the Yoneda algebra $B_n^!$. In this way we provided dictionaries relating modules over the Weyl algebras $A_n$ with graded modules over a finite dimensional selfinjective Koszul algebra $B_n^!$.

In this paper we will concentrate in the study of the enveloping algebra $U$ of the Lie algebra $s\ell(2,\mathbb{C})$ and its homogenized algebra $B$. In this case $B$ is a Koszul Artin-Schelter regular algebra of global dimension 4 and we can extend the methods used for the Weyl algebras to study the relations among the $U$-modules, the graded $B$ -modules, and the graded $B^!$-modules.

It is generally agreed that the classification of all modules over a wild finite dimensional algebra is impossible. It makes sense to look to families of indecomposable modules over a particular finite dimensional wild algebra, this is what we

---

*Date*: November 5, 2012.

2000 *Mathematics Subject Classification.* Primary 05C38, 15A15; Secondary 05A15, 15A18.

*Key words and phrases.* homogenized sl2, Koszul.

do in this paper for the selfinjective wild algebra $B^!$. By the standard methods of representation theory we do not know much about the $B^!$- modules, we only know by a result of [MZ], that the graded stable Auslander-Reiten components are all of type $ZA_\infty$. However, the algebra $B$ is the only Lie algebra for which all irreducible modules are known, in particular a family of modules $\{M(\lambda) \mid \lambda \in \mathbb{C}\}$ called the Verma modules is fully described. The so called weight modules and the related category $\mathcal{O}$ have been fully studied for $s\ell(2,\mathbb{C})$ [Ma]. We will here consider the corresponding categories of homogenized $B$-modules. Of particular interest is the homogenized Verma modules $V(\lambda)$ with $\lambda \in \mathbb{C}$, these modules are Koszul of global dimension 2 and we can describe the modules $W(\lambda)$ over the Yoneda algebra $B^!$corresponding to $V(\lambda)$ under Koszul duality. We prove that such modules $W(\lambda)$ belong to the mouth of a regular graded component, obtaining in this way an infinite number of components parametrized by $\lambda \in \mathbb{C}$.

## 2. The homogenized algebra $B$ of the enveloping algebra $U$ of $s\ell(2,\mathbb{C})$.

In this section we study the basic properties of the homogenized algebra $B$ of the enveloping algebra $U$ of $s\ell(2,\mathbb{C})$ and its Yoneda algebra $B^!$. We concentrate on the structure of these algebras, our main tool is the resolution given by Anick in [An]. The situation is very similar to the case of the Weyl algebras considered in [MMo1].

Through the paper $\mathbb{C}$ will denote the complex numbers, $s\ell(2,\mathbb{C})$ the $\mathbb{C}$-vector subspace of the space of two by two matrices $M_{2\times 2}(\mathbb{C})$ consisting of the matrices with zero trace. $s\ell(2,\mathbb{C})$ is a Lie algebra with bracket product $[X,Y] = XY - YX$. The enveloping algebra $U$ of $s\ell(2,\mathbb{C})$ is given by generators and relations by $U = \mathbb{C} < e, f, h > /L$, where $\mathbb{C} < e, f, h >$ is the free algebra with three generators: $e$, $f$, $h$ and $L$ is the ideal generated by the relations: $[e,f]$-$h$, $[h,e]$-$2e$, $[h,f]$+$2f$. It is well known that $U$ has a Poincare-Birkoff basis [Ma],[EW], this means that every element $u \in U$ can be written in a unique way as a combination $u = \sum\limits_{\ell} \sum\limits_{i+j+k=\ell} c_{i,j,k} e^i f^j h^k$ and $c_{i,j,k} \in \mathbb{C}$.

We will denote by $U$ the algebra defined by generators as: $B = \mathbb{C} < e, f, h, z > / I$ , where $\mathbb{C} < e, f, h, z >$ is the free algebra in four generators and $I$ is the ideal generated by the relations: $[e,f]$-$hz$, $[h,e]$-$2ez$, $[h,f]$+$2fz$, $[e,z]$, $[f,\ z]$, $[h,z]$. We will call $B$ the homogenized enveloping algebra of $s\ell(2,\mathbb{C})$. [Sm], [LbS].

We prove that $B$ has a Poincare-Birkoff basis, it is Koszul and it is Artin-Schelter regular.

We will use freely the definitions and theorems from [An].

We give to the generators $e$, $f$, $h$, $z$ the following order $z < f < e < h$ and to the words of $\mathbb{C} < e, f, h, z >$ the order grade lexicographic. There is a natural surjection $f : \mathbb{C} < e, f, h, z > \to B$.

Let $W$ be the monoid consisting of all words of $\mathbb{C} < e, f, h, z >$. The monoid $W$ is well ordered.

Given $u, v \in W$, following $[An]$, we will say that $v$ is a submonomial of $u$, $v < u$, if $v = 1$ or $v = x_{i_m} x_{i_{m-1}} ... x_{i_s}$ and $u = x_{i_1} x_{i_2} ... x_{i_t}$ , with $1 \leq m \leq s \leq t$ and $x_{i_j} \in W$.

Submonomial is a partial order on $W$.

A subset $M \subset W$ is an order ideal of monomials (o.i.m) if and only if $u \in M$ and $v < u$ implies $v \in M$.

Let $M$ be defined as $M = \{x \in W \mid f(x) \notin span f(y),\, y < x\}$.

We have from $[An]$:

**Lemma 1.** *The set $M$ is an o.i.m and the elements $f(x)$ for $x \in M$ form a basis of $B$ as $\mathbb{C}$-module.*

We compute in our case these basis.

The following diagrams illustrate the order of the monomials involved in the relations defining the ideal $I$.

$$\begin{tikzcd} & \cdot \arrow[rd, "z"] & \\ \cdot \arrow[ru, "h"] \arrow[r, "e"] \arrow[rd, "f"'] & \cdot \arrow[r, "f"] & \cdot \\ & \cdot \arrow[ru, "e"'] & \end{tikzcd}, \text{ where } fe < ef < hz$$

$$\begin{tikzcd} & \cdot \arrow[rd, "e"] & \\ \cdot \arrow[ru, "h"] \arrow[r, "e"] \arrow[rd, "e"'] & \cdot \arrow[r, "h"] & \cdot \\ & \cdot \arrow[ru, "z"'] & \end{tikzcd}, \text{ where } ez < eh < he$$

$$\begin{tikzcd} & \cdot \arrow[rd, "f"] & \\ \cdot \arrow[ru, "h"] \arrow[r, "f"] \arrow[rd, "f"'] & \cdot \arrow[r, "h"] & \cdot \\ & \cdot \arrow[ru, "z"'] & \end{tikzcd}, \text{ where } fz < fh < hf$$

$$\begin{tikzcd} & \cdot \arrow[rd, "z"] & \\ \cdot \arrow[ru, "e"] \arrow[rd, "z"'] & & \cdot \\ & \cdot \arrow[ru, "e"'] & \end{tikzcd}, \text{ where } ze < ez$$

$$\begin{tikzcd} & \cdot \arrow[rd, "z"] & \\ \cdot \arrow[ru, "f"] \arrow[rd, "z"'] & & \cdot \\ & \cdot \arrow[ru, "f"'] & \end{tikzcd}, \text{ where } zf < fz$$

$$\begin{tikzcd} & \cdot \arrow[rd, "z"] & \\ \cdot \arrow[ru, "h"] \arrow[rd, "z"'] & & \cdot \\ & \cdot \arrow[ru, "h"'] & \end{tikzcd}, \text{ where } zh < hz$$

It follows from the defining relations and the order we gave to the monomials that the paths: $hz$, $he$, $hf$, $ez$, $fz$ do not belong to $M$.

Moreover, $hz = ef - fe$ and $hz = zh$ , hence $ef - fe - zh = 0$ and $ef > fe$, $fe > zh$. Therefore: $ef \notin M$.

$M$ does not contain the paths $\{\ hz,\ he,\ hf,\ ez,\ fz,\ ef\}$.

It is clear that a word is in $M$ if and only if it is of the form $z^i f^j e^k h^\ell$.

As a consequence we have:

**Theorem 1.** *The algebra $B$ has a Poincare-Birkoff basis.*

Using the Poincare-Birkhoff basis we define a good filtration on $B$, as follows:

Let $b$ be an element of $B$. Then $b$ is written as $b = \sum\limits_{i,j,k,\ell \geq 0} c_{i,j,k,\ell} z^i f^j e^k h^\ell$, with $c_{i,j,k,\ell} \in \mathbb{C}$, such that all but a finite number of them are zero.

The degree of $b$ is the largest integer $n$ such that some $c_{i,j,k,\ell} \in \mathbb{C}$ with $j+k+\ell = n$ is non zero.

Let $\mathcal{F}_n$ be defined by $\mathcal{F}_n = \{\ b \in B \mid \text{degree(b)} \leq n\}$.

Then $\mathcal{F}_{-1} = 0$, $\mathcal{F}_0 = \mathbb{C}[z]$ , $\mathcal{F}_i \subset \mathcal{F}_{i+1}$, for all $i \geq 0$, $B = \bigcup\limits_{n \geq 0} \mathcal{F}_n$, $\mathcal{F}_n \mathcal{F}_m \subset \mathcal{F}_{n+m}$ and $B$ is generated as algebra by $\mathcal{F}_1$.

The associated graded algebra $\bigoplus\limits_{i \geq} \mathcal{F}_i / \mathcal{F}_{i-1}$ is isomorphic to the polynomial ring $\mathbb{C}[z, e, f, h]$.

Proceeding as in [MMo1], we can check this is a good filtration. We leave the details to the reader.

As a consequence of [Mi], we obtain:

**Theorem 2.** *The algebra $B$ is noetherian.*

Let $V_M$ be the obstruction set, $V_M = \{v \in W \mid v \notin M,\ u \lneqq v \text{ implies } u \in M\}$

It is clear that $V_M = \{\ hz,\ he,\ hf,\ ez,\ fz,\ ef\}$.

According to [An], the $0-$chains are precisely the arrows $V_0 = \{e,\ f,\ h,\ z\}$, the 1-chains $V_1 = V_M$, the 2-chains are the 2-overlaps $V_2$ ={efz, hfz, hez, hef }, the 3-chains are the 3-overlaps $V_3 =$ {hefz}.

There are no 4-chains.

According to Anick, there is a projective resolution of the only graded simple $S : *)$

$$0\to V_3 \otimes_{\mathbb{C}} B \to V_2 \otimes_{\mathbb{C}} B \to V_1 \otimes_{\mathbb{C}} B \to V_0 \otimes_{\mathbb{C}} B \to B \to S \to 0$$

which results minimal and linear, hence $B$ is a Koszul algebra of global dimension 4.

We will compute the Yoneda algebra $B^{\,!}$ of $B$.

We know by the general theory of Koszul algebras [GM1] ,[GM2], that $B^!$ is defined by quiver and relations as $B^! = \mathbb{C} < e, f, h, z >/\ I^{\perp}$ , where $\mathbb{C} < e, f, h, z >$ is the free algebra in four generators and $I^{\perp}$ is the ideal generated by the relations orthogonal to the relations defining $I$ with respect to the bilinear form defined in the paths of length two by $< \alpha\beta, \alpha'\beta' >= \begin{cases} 1 & \text{if } \alpha = \alpha',\ \beta = \beta' \\ 0 & \text{otherwise} \end{cases}$

Let's compute $I^{\perp}$.

$< e^2,\ ef - fe - hz>=< f^2,\ ef - fe - hz>=< h^2,\ ef - fe - hz>=< z^2,\ ef - fe - hz>$=0'.

$< e^2,\ he - eh - 2ez>=< f^2,\ he - eh - 2ez>=< h^2,\ he - eh - 2ez>=< z^2,\ he - eh - 2ez>$=0

$< e^2,\ hf - fh + 2fz>=< f^2,\ hf - fh + 2fz>=< h^2,\ hf - fh + 2fz>=< z^2,\ hf - fh + 2fz>$=0

$< e^2,\ fz - zf>=< f^2,\ fz - zf>=< h^2,\ fz - zf>=< z^2,\ fz - zf>$=0

$< e^2,\ ez - ze>=< f^2,\ ez - ze>=< h^2,\ ez - ze>=< z^2,\ ez - ze>$=0

$< e^2,\ hz - zh>=< f^2,\ hz - zh>=< h^2,\ hz - zh>=< z^2,\ hz - zh>$=0.

It follows $e^2,\ f^2,\ h^2,\ z^2 \in I^{\perp}$.

$< ef+fe, ef-fe-hz>=< ef+fe, he-eh-2ez>=< ef+fe, hf-fh+2fz>=< ef+fe,\ fz-zf>=< ef+fe,\ ez-ze>=< ef+fe,\ hz-zh>$=0

$< he+eh, ef-fe-hz>=< he+eh, he-eh-2ez>=< he+eh, hf-fh+2fz>=< he+eh,\ fz-zf>=< he+eh,\ ez-ze>=< he+eh,\ hz-zh>$=0

$< hf+fh, ef-fe-hz>=< hf+fh, he-eh-2ez>=< hf+fh, hf-fh+2fz>=< hf+fh,\ fz-zf>=< hf+fh,\ ez-ze>=< hf+fh,\ hz-zh>$=0.

Therefore: $ef + fe,\ he + eh,\ hf + fh \in I^{\perp}$.

$< hz+zh+ef, ef-fe-hz>=< hz+zh+ef\ , he-eh-2ez>=< hz+zh+ef\ , hf-fh+2fz>=< hz+zh+ef\ , fz-zf>=< hz+zh+ef\ , ez-ze>=< hz+zh+ef,\ hz-zh>$=0

$< ez+ze-2eh, ef-fe-hz>=<ez+ze-2eh\ , he-eh-2ez>=<ez+ze-2eh\ , hf-fh+2fz>=<ez+ze-2eh\ , fz-zf>=<ez+ze-2eh\ , ez-ze>=<ez+ze-2eh\ , hz-zh>$=0

$< fz+zf-2hf, ef-fe-hz>=<fz+zf-2hf\ , he-eh-2ez>=<fz+zf-2hf, hf-fh+2fz>=<\ , fz-zf>=< fz+zf-2hf, ez-ze>=<\ , hz-zh>$=0

In consequence, $hz + zh + ef,\ ez + ze - 2eh,\ fz + zf - 2hf \in I^{\perp}$.

We will use the following notation: $(a, b) = ab+ba$ and $S$=$\{\ e^2,\ f^2,\ h^2,\ z^2,\ (e, f), (e, h),\ (f, h),\ (h, z) + ef,\ (e, z) - 2eh,\ (f, z) - 2hf\}$.

We have proved the inclusion $S \subset I^{\perp}$.We will check that $S$ generates $I^{\perp}$.

Let $b$ be an element of degree two in $I^{\perp}$. Then $b$ has the following form:

$b$= $a_1e^2$+ $a_2$ $f^2+a_3h^2$+ $a_4$ $z^2+a_{12}fh+a_{21}hf+a_{13}fe+a_{31}ef+a_{14}fz+a_{41}zf+a_{23}he+a_{32}eh+a_{24}hz+a_{42}zh+a_{34}ez+a_{43}ze$.

which can be rewritten as:

$b = a_1e^2+ a_2\ f^2 + a_3h^2+ a_4\ z^2 + a_{12}(fh+hf)+a_{13}(fe+ef)+a_{14}(fz+zf-2hf)+a_{23}(he+eh)+a_{24}(hz+zh+ef)+a_{34}(ez+ze-2eh)+b'$.

Where $b' = (a_{21}-a_{12}+2a_{14})hf+(a_{31}-a_{13}-a_{24})ef+(a_{32}-a_{23}+2a_{34})eh+(a_{41}-a_{14})zf+(a_{42}-a_{24})zh+(a_{43}-a_{34})ze$.

It follows $b' \in I^{\perp}$.

Then $< b',fz-zf >= a_{14}-a_{41} = 0$, $< b',hz-zh >= a_{24}-a_{42} = 0$, $< b',ez-ze >= a_{34}-a_{43} = 0$.

$< b',\ hf-fh-2fz >= 0 = a_{21}-a_{12}+2a_{14}$, $< b',\ ef-fe-hz >= 0 = a_{31}-a_{13}-a_{24}$, $< b',he-eh-2ez >= -((a_{32}-a_{23}+2a_{34}) = 0$.

We have proved $b' = 0$ and $I^{\perp}$ is generated by $S$, this is: { $e^2$, $f^2$, $h^2$, $z^2$, $(e,f)$, $(e,h)$, $(f,h)$, $(h,z)+ef$, $(e,z)-2eh$, $(f,z)-2hf$} $= I^{\perp}$.

The algebra $B^!$ is graded and $B^! = L_0 \oplus L_1 \oplus L_2 \oplus L_3 \oplus L_4$, with $L_0 = \mathbb{C}1$, $L_1 = \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}z$, $L_2$ generated by { $fe$, $he$, $fh$, $fz$, $hz$, $ez$}, $L_3$ generated by {$fhe$, $fez$, $fhz$, $hez$} and $L_4$ generated by { $fhz$}.

We most prove $dim\ _{\mathbb{C}}L_2 = 6$, dim $_{\mathbb{C}}L_3 = 4$, dim $_{\mathbb{C}}L_4 = 1$.

Using the resolution for the simple graded $B$-module $S$ *) and the Koszul property [GM1] we get what we claimed, $dim\ _{\mathbb{C}}L_2 = 6$, $dim\ _{\mathbb{C}}L_3 = 4$, dim $_{\mathbb{C}}L_4 = 1$.

It follows that the algebra $B^!$ has simple socle, hence it is selfinjective [Y].

By Koszul theory, [Sm] the algebra $B$ is Artin-Schelter regular.

It will be clear from the relations that the algebra $B^!$ has a structure related with the exterior algebra.

Let $C^!$ be the exterior algebra: $C^!$=$\mathbb{C}$<$e,f,h$>/<$e^2,f^2,h^2,(e,f),(e,h),(f,z)$>. Then $C^!$ is a subalgebra of $B^!$ and $B^!$ decomposes $B^! = C^! \oplus C^!z$ as a left and $B^! == C^! \oplus zC^!$ as a right $C^!$ -module .

It is clear that $B/zB \cong C$ where $C$ is the polynomial algebra $\mathbb{C}[e.f,h]$.

Observe that as in [MMo1] $C$ appears as a quotient of $B$ and $C^!$ as a subalgebra of $B^!$.

In the next proposition we describe the relations between the homogenized enveloping algebra $B$ and the usual enveloping algebra of $s\ell(2,\mathbb{C})$.

**Proposition 1.** *There is an isomorphism of* $\mathbb{C}$*-algebras* $B/(z-1)B \cong U$.

*Proof.* We have $\mathbb{C}$-algebra homomorphisms $\mathbb{C} < e,f,h > \xrightarrow{j} B \xrightarrow{\pi} B/(z-1)B$. If $q(e,f,h)$ is a word in $e$, $f$, $h$ and $j(q(e,f,h)) = q(e,f,h)+I$.

Let $\varphi$ be the composition $\varphi = \pi j$. Then $\varphi(he-eh-2e) = j((he-eh-2ez)+2e(z-1))+(z-1)B = 0$.

Similarly, $\varphi(ef-fe-h) = 0$ and $\varphi(hf-fh+2f) = 0$.

The ideal $L =< ef-fe-h,\ he-eh-2e,\ hf-fh+2f >$ is contained in the kernel of $\varphi$ and $\varphi$ induces an homomorphism $\overline{\varphi}: U \to B/(z-1)B$.

We prove that $\overline{\varphi}$ is an isomorphism.

i) $\overline{\varphi}$ is injective.

Let $u$ be an element of $U$ and $u = \sum c_{i,j.k}e^if^jh^k$ with $c_{i,j.k}$ a finite number of elements in $\mathbb{C}$. $\overline{\varphi}$ $(u)$=0 if and only if $\sum c_{i,j.k}e^if^jh^k = b(z-1) \in (z-1)B$ and

$b$ can be expressed as $b = q_0 + q_1 z + q_2 z^2 + ...q_t z^t$. We write $\sum c_{i,j.k} e^i f^j h^k = -q_0 + (q_0 - q_1)z + (q_1 - q_2)z^2 + ...(q_{t-1} - q_t)z^t + q_t z^{t+1}$.

Since $e$, $f$, $h$, $z$ induce a Poincare-Birkoff basis in $B$, $\sum c_{i,j.k} e^i f^j h^k = -q_0$ and $q_0 = q_1 = ... = q_t = 0$.

ii) $\overline{\varphi}$ is surjective.

Let $b + (z-1)B$ be an element of $B/(z-1)B$ and $b = q_0 + q_1 z + q_2 z^2 + ...q_t z^t$.

We write $z^k = ((z-1)+1)^k = (z-1)p(z) + 1$ with $p(z)$a polynomial in $z$.

Then $b + (z-1)B = q_0 + q_1 + q_2 + ...q_t + (z-1)B$.

Therefore: $\overline{\varphi}(q_0 + q_1 + q_2 + ...q_t) = b + (z-1)B$. □

We give next another characterization of $U$.

We will follow very closely the ideas and results form [MMo1], although the next propositions were given in [MMo1], we include them here for completeness.

By construction $B$ is a $\mathbb{C}[z]$-algebra.

Consider the multiplicative subset $S = \{1, z, z^2,..\}$ of $\mathbb{C}[z]$ and denote by $\mathbb{C}[z]_S$ the graded localization. It was proved in [MMo1] that there is a $\mathbb{C}$-algebras isomorphism $\mathbb{C}[z]_S \cong \mathbb{C}[z, z^{-1}]$, where $\mathbb{C}[z, z^{-1}]$ denotes the Laurent polynomials.

The algebra $B_z = B \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}]$ is a $\mathbb{Z}$-graded algebra with homogeneous elements $b/z^k$, where b is homogeneous and the degree of $b/z^k$ is deg($b/z^k$)=deg(b)-k .

There exists a homomorphism of graded algebras $\varphi : B \to B_z$ given by $\varphi(b) = b/1 = b \otimes 1$. Consider the composition of rings homomorphisms: $B \overset{\varphi}{\to} B_z \overset{\pi}{\to} B_z/(z-1)B_z$.Clearly, $(z-1)B \subset \ker \pi\varphi$.

Let $b$ be an element of $B$ such that $b/1 \in (z-1)B_z$.This is: $b/1 = (z-1)b'/z^k$.

Therefore: $z^\ell b = (z-1)z^t b' = (z-1)p(z)b + b$ and $p(z)$ a polynomial in $z$. Hence, $b = (z-1)b'' \in (z-1)B$.

We have a factorization of $\pi\varphi$:

$$\begin{tikzcd} B \arrow[r, "\varphi"] \arrow[dr, "q"'] & B_z \arrow[r, "\pi"] & B_z/(z-1)B_z \\ & B/(z-1)B \arrow[ur, "\psi"'] & \end{tikzcd}$$

with $\psi$ an injective homomorphism of $\mathbb{C}$-algebras.

Let $b/z^k + (z-1)B_z$ be an element of $B_z/(z-1)B_z$. The element $b/z^k$ can be written as: $b/z^k = b_\ell/z^\ell + b_{\ell-1}/z^{\ell-1} + ...b_0 + b_1 z + ...b_m z^m$, where each $b_i$ is a polynomial in $e$, $f$, $h$.

For each $i$, $b_i/1 = z^i b_i/ z^i = (z-1)p(z)b_i/z^i + b_i/z^i$. Then $b_i/z^i + (z-1)B_z = b_i/1 + (z-1)B_z$.

It follows $b/z^k + (z-1)B_z = (b_\ell + b_{\ell-1} + ...b_0 + b_1 + ...b_m)/1 + (z-1)B_z$.

The element $u = b_\ell + b_{\ell-1} + ...b_0 + b_1 + ...b_m$ is a polynomial in $e$, $f$, $h$ , therefore it is an element of $U = B/(z-1)B$ and $\psi(u) = b/z^k + (z-1)B_z$.

We have proved $\psi$ is an isomorphism with inverse $\psi^{-1}$. The composition $\tau = \psi^{-1}\pi$ makes the diagram:

$$\begin{tikzcd} B \arrow[rr, "\varphi"] \arrow[dr, "q"'] & & B_z \arrow[dl, "\tau"] \\ & B/(z-1)B & \end{tikzcd}$$

commute.

The algebra $B_z$ is $\mathbb{Z}$-graded and has in degree zero a subalgebra $(B_z)_0$. There is an inclusion $(B_z)_0 \to B_z$ and a projection $B_z \to B_z/(z-1)B_z$. Denote by $\theta :(B_z)_0 \to B_z/(z-1)B_z$ the composition.

**Proposition 2.** *The morphism* $\theta:(B_z)_0 \to B_z/(z-1)B_z$ *is an isomorphism.*

*Proof.* Let $u = \sum_{i=0}^{m} g_i(f,\, h,\, e) z^{-n_i}$ be an element of $(B_z)_0$, this is: each $g_i(f,\, h,\, e)$ is a polynomial in $e$, $f$, $h$ of degree $n_i$ and $n_0 > n_1 > ... > n_m$.

We can rewrite $u$ as $u = z^{-n_0-1}\sum_{i=0}^{m} g_i(f,\, h,\, e) z^{n_0+1-n_i} =$

$z^{-n_0-1}(\sum_{i=0}^{m}(g_i(f,\, h,\, e) + (z-1)p_i(z)g_i(f,\, h,\, e))$.

We write $g(\, f,\, h,\, e) = \sum_{i=0}^{m}(g_i(f,\, h,\, e)$.

Hence; $\theta(u) = u + (z-1)B_z = z^{-n_0-1}g(f,\, h,\, e) + (z-1)B_z$.

Assume $\theta(u) = 0$. Then $z^{-n_0-1}g(f,\, h,\, e) = (z-1)b'z^{-k}$.

It follows that there exist integers $s, t \geq 0$ with

$z^t g(f,\, h,\, e) = g(f,h,e) + (z-1)p(z)g(f,h,e) = (z-1)b''z^s$.

Therefore: $g(f,\, h,\, e) = (z-1)\overline{b}$ with

$\overline{b} = b_0(f,\, h,\, e) + b_1(f,\, h,\, e)z + b_2(f,\, h,\, e)z^2 + ...b_k(f,\, h,\, e)z^k$.

As above,

$g(f,\, h,\, e) = -b_0(f,\, h,\, e) + (b_0(f,\, h,\, e) - b_1(f,\, h,\, e))z + (b_1(f,\, h,\, e) - b_2(f,\, h,\, e))z^2 + ...(b_{k-1}(f,\, h,\, e) - b_k(f,\, h,\, e))z^k + b_k(f,\, h,\, e)z^{k+1}$.

It follows $g(f,\, h,\, e) = -b_0(f,\, h,\, e)$ and $b_0(f,\, h,\, e) = b_1(f,\, h,\, e)) = b_2(f,\, h,\, e) = ... = b_k(f,\, h,\, e) = 0$.

We have proved $\sum_{i=0}^{m}(g_i(f,\, h,\, e)$=0 with each $g_i(f,\, h,\, e)$ homogeneous of degree $n_i$.

It follows $g_i(f,\, h,\, e) = 0$ for all $i$ and $u = 0$.

We prove next that $\theta$ is surjective.

Let $b/z^k + (z-1)B_z$ be an element of $B_z/(z-1)B_z$, $b = \sum_{i=0}^{m} b_i$ and $b_i$ homogeneous components of degree $n_i$ with $n_0 < n_1 < ... < n_m$.

As before, $z^{n_m-n_i}b_i = (z-1)p_i(z)b_i + b_i$, the element of $B$, $b'_i = z^{n_m-n_i}b_i$ is homogeneous of degree $n_m$ and $b' = \sum_{i=0}^{m} b'_i$ is homogeneous of degree $n_m = \ell$.

Then $b/z^k + (z-1)B_z = b'/z^k + (z-1)B_z$.

If $\ell > k$, then $b'/z^k + (z-1)B_z = z^{\ell-k}b'/z^\ell + (z-1)B_z = b'/z^\ell + (z-1)B_z$ with degree $b'/z^\ell = 0$.

If $\ell < k$, then $b''/z^k = z^{k-\ell}b'/z^k = b'/z^k + (z-1)p(z)b'/z^k$, where $b'' = z^{k-\ell}b$ has degree $k$.

$b''/z^k + (z-1)B_z = b'/z^k + (z-1)B_z = b/z^k + (z-1)B_z$.

In the first case $\theta(b'/z^\ell) = b/z^k + (z-1)B_z$.

In the second case $\theta(b''/z^k) = b/z^k + (z-1)B_z$.

We have proved $\theta$ is an isomorphism. □

With the identification $U \cong B/(z-1)B \cong B_z/(z-1)B_z$ we have proved $B_z$ has a $U$ in degree zero.

**Theorem 3.** *There exist an isomorphism of graded rings:* $U[z, z^{-1}] = U \otimes_{\mathbb{C}} \mathbb{C}[z, z^{-1}] \cong B \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}]$.

*Proof.* Identifying $U$ with $(B_z)_0$ we will prove that the morphism:

$\mu : (B_z)_0 \otimes_{\mathbb{C}} \mathbb{C}[z, z^{-1}] \to B_z$ given by $\mu(bz^{-\ell} \otimes z^k) = bz^{k-\ell}$ is a ring isomorphism.

Let $b/z^\ell$ be an homogeneous element of $B_z$ with degree$b = \ell + k$.

Then $b/z^\ell = z^k b/z^{\ell+k}$ and $\mu(b/z^{\ell+k} \otimes z^k) = b/z^\ell$ and $\mu$ is surjective.

Let $b/z^k$ be an homogeneous element of degree zero.

Then $b = \sum_{i=0}^{k} g_i(f,\, h,\, e)z^{k-i}$ with each $g_i(f,\, h,\, e)$ a polynomial in $f$, $e$, $h$ which is either zero, or degree$(g_i(f,\, h,\, e)) = i$.

$\mu(b/z^k \otimes z^\ell) = z^{\ell-k} \sum_{i=0}^{k} g_i(f,\, h,\, e)z^{k-i} = 0$ in $B_z$.

There exist $t \geq 0$ such that $z^t \sum_{i=0}^{k} g_i(f,\, h,\, e)z^{k-i} = 0$ in $B.z$

Then $b/z^k \otimes z^\ell = z^t b/z^{k+t} \otimes z^\ell = 0$.

We have proved $\mu$ is an isomorphism. □

The ring $U[z, z^{-1}]$ is a strongly $\mathbb{Z}$-graded. The ring homomorphism $U \to U[z, z^{-1}]$ induces functors:

$U[z, z^{-1}] \otimes_U - : Mod_U \to Gr_{U[z,z^{-1}]}$ and $res_U : Gr_{U[z,z^{-1}]} \to Mod_U$.

By Dade's theorem [Da] we have:

**Theorem 4.** *The functors $U[z, z^{-1}] \otimes_U -$ and $res_U$ are inverse exact equivalences.*

**Corollary 1.** *The equivalences $U[z, z^{-1}] \otimes_U -$ and $res_U$ preserve projective and irreducible modules and send left ideals to left ideals giving an order preserving bijection.*

**Definition 1.** *Given a $B$-module $M$ we define the $z$-torsion of $M$ as $t_z(M) = \{m \in M \mid \text{there exists } n \geq 0 \text{ with } z^n m = 0\}$. The module $M$ is of $z$-torsion if $t_z(M) = M$ and $z$-torsion free if $t_z(M) = 0$.*

$t_z(M)$ is a submodule of $M$ and a map $f : M \to N$ restricts to a map $f_{|t_z(M)} : t_z(M) \to t_z(N)$ in this way $t_z(-)$ is a subfunctor of the identity with $t_z(t_z(M)) = t_z(M)$,

For any $B$-module $M$ there is an exact sequence:

$$0 \to \quad t_z(M) \quad \to \quad M \quad \to \quad M/t_z(M) \quad \to 0$$

with $t_z(M)$ of $z$-torsion and $M/t_z(M)$ $z$-torsion free.

The kernel of the natural morphism $M \to M_z = B_z \otimes_B M$ is $t_z(M)$.

In the next proposition we describe as a particular case known facts concerning any localization [P].

**Proposition 3.** *The following statements hold:*

*i) Given a graded map of $B$-modules $f : M \to N$ the map induced in the localization: $f_z : M_z \to N_z$ is zero if and only if $f$ factors through a $z$-torsion module.*

*ii) Assume the localized modules $M_z$ is finitely generated and let $\varphi : M_z \to N_z$ be a morphism of graded $B_z$-modules. Then there exists an integer $k \geq 0$ and a morphism of $B$-modules $f : z^k M \to N$ such that the composition $M_z \xrightarrow{\sigma} (z^k M)_z \xrightarrow{f_z} N_z$, $f_z\sigma = \varphi$ and $\sigma$ is an isomorphism of graded $B_z$-modules.*

*iii) Let $M$ be a finitely generated $B_z$-module. Then there exists a finitely generated graded $B$-submodule $\overline{M}$ of $M$ such that $\overline{M}_z \cong M_z$.*

*Proof.* i) Let $f : M \to N$ be a morphism of graded $B$-modules with $f_z = 0$ and $m \in M$. Then $f(m)/1 = 0$ implies there exists an integer $k \geq 0$ with $z^k f(m) = 0$, hence $f(M)$ is of $z$-torsion and $f$ factors $M \to f(M) \to N$.

Conversely, if $f = hg$ with $g : M \to L$ and $h : L \to N$ maps with $L$ of $z$-torsion, the $L_z = 0$ and $f_z = h_z g_z = 0$.

ii) Let $\varphi : M_z \to N_z$ be a morphism of graded $B_z$-module and $M_z$, $N_z$ finitely generated. Assume $m_1$, $m_2$, ... $m_k$ generate $M_z$ and deg$ree$ $m_i = d_i$ and let $d$ be $d = \max\{d_i\}$

$\varphi(m_j) = \sum n_{ij} \otimes z^{k_{ij}}$ with degree $n_{ij} + k_{ij} = d_i$.

If $k_{ij} > 0$,then $n_{ij} \otimes z^{k_{ij}} = n_{ij}z^{k_{ij}} \otimes 1$ and we may assume that in the expression $\varphi(m_j) = \sum n_{ij} \otimes z^{k_{ij}}$ all $k_{ij} \leq 0$. Let $k$ be $k = \max\{-k_{ij}\}$. Then $k + k_{ij} \geq 0$. Hence; $\varphi(m_j) = \sum n_{ij} z^{k+k_{ij}} \otimes z^{-k} = n_j \otimes z^{-k}$ and degree $n_j = k + d_j$.

Let $j : M \to M_z = M \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}]$ is the localization map $j(m) = m/1 = m \otimes 1$ and $\varphi : M_z = M \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}] \to N_z = N \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}]$ is the given map.

Let $f' : z^k M \to N \otimes_{\mathbb{C}[z]} \mathbb{C}[z, z^{-1}]$ be the map $f(z^k m_j) = z^k n_j \otimes z^{-k} = n_j \otimes 1$ and $f'(z^k M) \subset N \otimes 1 \cong N$ restricting the image and identifying $N \otimes 1 = N$ we obtain a map $f : z^k M \to N$ given by $f(z^k m_j) = n_j$.

There is an exact sequence $0 \to z^k M \xrightarrow{i} M \to M/z^k M \to 0$, where $M/z^k M$ is of $z$-torsion. Localizing we obtain an exact sequence: $0 \to (\ z^k M)_z \xrightarrow{i_z} M_z \to (M/z^k M)_z \to 0$ with $(M/z^k M)_z = 0$ and the map $\sigma = i_z$ is an isomorphism.

The morphism $f_z$ is defined as $f_z(m/z^\ell) = f_z(z^k(m/z^{k+\ell}) = f(z^k m)/z^{k+\ell} = \varphi(z^k m)/z^{k+\ell} = z^k \varphi(m)/z^{k+\ell} = \varphi(m)/z^\ell = \varphi(m/z^\ell)$.

iii) Let $M$ be a finitely generated $B_z$-module with homogeneous generators $m_1$, $m_2$,... $m_k$ of degree $m_i = d_i$.

By restriction, $M$ is a $B$-module. Let $\overline{M}$ be the $B$-submodule of $M$ generated by $m_1$, $m_2$,... $m_k$.

We need to check that under localization $\overline{M}\,_z$ is isomorphic to $M$ as $B_z$-modules.

Localizing, $\overline{M}\,_z \cong B_z \otimes_B \overline{M} \cong \mathbb{C}[z, z^{-1}] \otimes_{\mathbb{C}[z]} B \otimes_B \overline{M} \cong \mathbb{C}[z, z^{-1}] \otimes_{\mathbb{C}[z]} \overline{M}$. The homogeneous elements of $\overline{M}\,_z$ are of the form $z^{-k} \otimes m$, let $\mu : \mathbb{C}[z, z^{-1}] \otimes_{\mathbb{C}[z]} \overline{M} \to M$ be the multiplication map $\mu(z^{-k} \otimes m) = z^{-k} m$, hence $\mu(z^{-k} \otimes m) = 0 = z^{-k} m$ implies $z^k z^{-k} m = m = 0$.

and $\mu$ is injective.

Let $m$ be an homogeneous element of $M$ of degree $k$. Then $m = \sum b_i / z^{n_i} m_i$ and degree $b_i + d_i - n_i = k$. Let $n$ be $n = \max\{n_i\}$.

Therefore: $z^n m = \sum z^{n-n_i} b_i / m = \overline{m}$ and $\overline{m} \in \overline{M}$ with degree$\overline{m} = k + n$ .

It follows, $\mu(z^{-n} \otimes \overline{m}) = m$. $\square$

**Corollary 2.** *Let $M$ and $N$ be graded $B$-modules with $M$ finitely generated. A morphism $\varphi : M_z \to N_z$ is an isomorphism if and only if the map $f : z^k M \to N$ such that $f_z = \varphi$ has kernel and cokernel of $z$-torsion.*

## 3. The homogenized Verma modules

In the previous section we saw that there is an exact equivalence between the category of graded $B_z$-module, $Gr_{B_z}$ and the category of $U$-modules, $Mod_U$,where $U$ is the enveloping algebra of $s\ell(2,\mathbb{C})$, $B$ its homogenization and $B_z$ is the localization of $B$ in $z$. We also sketched the relations between the category of graded $B$-modules and the category of graded $B_z$-modules. We remarked that the algebra $B$ is Koszul and studied both the structure of $B$ and the structure of its Yoneda algebra $B^!$. In this setting families of modules over the enveloping algebra are know and we know that $B^!$ is a selfinjective Koszul wild algebra whose graded stable Auslander-Reiten components are of type $ZA_\infty$ [MZ]. The aim of the section is to lift known categories of $U$- modules to categories of $B$-modules. What we mean is; to describe categories of $B$-modules such that, when we apply the localization functor and restrict them to the degree zero part of $U$ of $B_z$, we obtain the given categories of $U$-modules. In some cases we will be able to use Koszul duality to obtain families of $B^!$-modules and see how they are distributed among the Auslander-Reiten components. Of special interest are the Verma modules, we study homogenized versions of Verma $B$-modules and of the so called, category $\mathcal{O}$ [Ma] .

3.1. **The Verma $B$-modules.** For each $\lambda \in \mathbb{C}$ we define the homogenized left ideal $I_\lambda$ of $B$ by $I_\lambda = Be + B(h - \lambda z)$ and the homogenized Verma module $V(\lambda) = B/I_\lambda$.

An homogeneous element $\overline{b}$ of $V(\lambda)$ is of the form:

$\overline{b} = b + I_\lambda$ with $b = \sum\limits_{i+\ j+m+\ell=k} c_{i,\ j,m,\ell} f^i h^j e^m z^\ell$ which can be written as:

$$b = \sum_{\substack{i+\ j+m+\ell=k \\ m\geq 1}} c_{i,\ j,m,\ell}\ z^\ell f^i h^j e^m + \sum_{i+\ j+\ell=k} c'_{i,\ j,\ell} z^\ell f^i h^j \text{ and}$$

$$\overline{b} = \sum_{i+\ j+\ell=k} c'_{i,\ j,\ell} z^\ell f^i (h - \lambda z) + \lambda z)^j + I_\lambda =$$

$$\sum_{t=0}^{\ell} \sum_{i+\ j+\ell=k} c'_{i,\ j,\ell} z^\ell f^i (h - \lambda z)^{j-t} (\lambda z)^t + I_\lambda = \sum_{i+\ j+\ell=k} c'_{i,\ j,\ell} \lambda^j z^{\ell+j} f^i + I_\lambda.$$

It follows that the monomials $\{\ f^i z^m\}$ generate $V(\lambda)$ as $\mathbb{C}$-vector space. We prove they form a basis.

We will prove by induction on $k$ that $\sum\limits_{i+j=k} a_{ij} f^i z^j \in I_\lambda$ implies $a_{ij} = 0$.

For $k = 1$, $a_{01} z + a_{10} f = b_1 e + b_2(h - \lambda z)$ implies $b_1 = 0$ and $b_2 = 0$, therefore: $a_{01} = a_{10} = 0$.

Assume $k > 1$.

If $\sum\limits_{i+j=k} a_{ij} f^i z^j \in I_\lambda$, then $\sum\limits_{i+j=k} a_{ij} f^i z^j =$

$$\sum_{i+\ j+m+\ell=k-1} c_{i,\ j,m,\ell}\ z^\ell f^i h^j e^{m+1} + \sum_{i+\ j+m+\ell=k-1} c'_{i,\ j,\ell} z^\ell f^i h^j e^m (h - \lambda z).$$

Since $eh = he - 2ez$, it follows by induction that $e^m h = he^m - 2me^m z$ and $e^m(h - \lambda z) = he^m - (2m + \lambda)e^m z$.

Then $\sum\limits_{i+j=k} a_{ij} f^i z^j =$

$$\sum_{i+\ j+m+\ell=k-1} c''_{i,\ j,m,\ell}\ z^\ell f^i h^j e^{m+1} + \sum_{i+\ j+\ell=k-1} d_{i,\ j,\ell} z^\ell f^i h^j (h - \lambda z).$$

This implies $c''_{i,\ j,m,\ell} = 0$ .

Hence we have equalities:

$$\sum_{i+j=k} a_{ij} f^i z^j = z(\sum_{i+\ j+\ell=k-1} d'_{i,\ j,\ell} z^{\ell-1} f^i h^j (h - \lambda z)) + \sum_{i+\ j=k-1} d''_{i,j} f^i h^j (h - \lambda z)$$

$$= z\left( (\sum_{i+\ j+\ell=k-1} d'_{i,\ j,\ell} z^{\ell-1} f^i h^j (h - \lambda z)) - \lambda \sum_{i+\ j=k-1} d''_{i,j} f^i h^j \right) + \sum_{i+\ j=k-1} d''_{i,j} f^i h^{j+1}.$$

Therefore: $d''_{i,j} = 0$, which implies $a_{k,0} = 0$

In the equality: $\sum\limits_{i+j=k} a_{ij}f^iz^j = z \ (\sum\limits_{i+\ j+\ell=k-1} d'_{i,\ j,\ell}z^{\ell-1}f^ih^j(h-\lambda z))$ we can cancel $z$ in both sides to have:

$$\sum_{i+j=k-1} a_{ij}f^iz^j = \sum_{i+\ j+\ell=k-1} d'_{i,\ j,\ell}z^{\ell-1}f^ih^j(h-\lambda z).$$

It follows by induction $a_{ij}=0$ for all $i$, $j$.

We have proved the following:

**Proposition 4.** *For each $\lambda \in \mathbb{C}$ the monomials $\{\ f^iz^m\}$ form a $\mathbb{C}$- basis of the homogenized Verma module $V(\lambda) = B/I_\lambda$, where $I_\lambda = Be + B(h-\lambda z)$.*

**Corollary 3.** *The homogenized Verma module $V(\lambda) = B/I_\lambda$ is $z$-torsion free.*

The exact sequence: $0 \to I_\lambda \to B \to B/I_\lambda \to 0$ induces by localization an exact sequence of $B_z$-modules $0 \to (I_\lambda)_z \to B_z \to (B/I_\lambda)_z \to 0$, restricting to the degree zero part we get an exact sequence:

$$0 \to ((I_\lambda)_z)_0 \to (B_z)_0 \to ((B/I_\lambda)_z)_0 \to 0$$

We proved the isomorphisms: $U \cong B/(z-1)B \cong B_z/(z-1)B_z \cong (B_z)_0$.

In the other hand, $(I_\lambda)_z = B_z \otimes_B I_\lambda = B_z \otimes_B Be + B_z \otimes_B B(h-\lambda z) = B_ze + B_z(h-\lambda z) = B_ze/z + B_z(h/z-\lambda)$.

Hence; $((I_\lambda)_z)_0 = (B_z)_0e/z + (B_z)_0(h/z-\lambda) \cong (B_z/(z-1)B_z)e/z + (B_z/(z-1)B_z)(h/z-\lambda)$..

As above, $e+(z-1)B_z = e/z+(z-1)B_z$ and $h-\lambda z+(z-1)B_z = h-\lambda+(z-1)B_z$.

It follows $((I_\lambda)_z)_0 = (B_z/(z-1)B_z)e + (B_z/(z-1)B_z)(h-\lambda) \cong Ue + U(h-\lambda)$.

Using these isomorphism we obtain an exact sequence:

$0 \to Ue + U(h-\lambda) \to U \to ((B/I_\lambda)_z)_0 \to 0$.

Therefore: $(V(\lambda))_0$ is the usual Verma $U$-module which we will denote by $M(\lambda)$.

It is well known [Ma], which Verma $U$-modules $M(\lambda)$ are irreducible.

**Proposition 5.** *A Verma $U$-modules $M(\lambda)$ is irreducible if and only if $\lambda \notin \mathbb{N} \cup \{0\}$.*

*If $n \in \mathbb{N} \cup \{0\}$, then the Verma $U$-modules $M(n)$ is indecomposable. Furthermore, the module $M(-n-2)$ is the unique simple submodule of $M(n)$ and $M(n)/M(-n-2){=}V^{(n+1)}$ is the unique finite dimensional simple of dimension $n+1$.*

As a consequence of this proposition and the exact equivalences:

$res_U : Gr_{B_z} = Gr_{U[z,z^{-1}]} \to Mod_U$ and $U[z,z^{-1}] \otimes_U - : Mod_U \to Gr_{B_z}$ it follows:

**Proposition 6.** *The localization of the homogenized Verma $B$-module $V(\lambda)_z$ is irreducible if and only if $\lambda \notin \mathbb{N} \cup \{0\}$.*

From this proposition we obtain the following properties of the homogenized Verma modules.

**Proposition 7.** *i) Given a non zero submodule $X$ of the homogenized Verma $B$-module $V(\lambda)$ with $\lambda \notin \mathbb{N} \cup \{0\}$ the module $V(\lambda)/X$ is $z$-torsion.*

*ii) The module $V(\lambda)$ is indecomposable for any $\lambda \in C$.*

*Proof.* i) Is a consequence of the previous proposition and ii) follows from the fact that $V(\lambda)$ has an indecomposable graded projective cover. □

As a consequence of Proposition 5, we have:

**Proposition 8.** *For $n \in \mathbb{N}\cup\{0\}$the homogenized Verma module $V(n)$ induces exact sequences:*

*$0 \to V(-n-2) \to V(n) \to V(n)/V(-n-2) \to 0$ and $0 \to V(-n-2)_z \to V(n)_z \to (V(n)/V(-n-2))_z \to 0$ with $V(-n-2)_z$ the unique simple submodule of $V(n)_z$ and $(V(n)/V(-n-2))_z$ has in degree zero the simple $U$-module of dimension $n+1$, $V^{(n+1)}$.*

In the next theorem we prove that the homogenized Verma modules are Koszul of global dimension two.

**Theorem 5.** *Let $V(\lambda)$ be a homogenized Verma $B$-module. Then $V(\lambda)$ has a minimal projective resolution: $0 \to B[-2] \overset{d_2}{\to} B \oplus B[-1] \overset{d_1}{\to} B \to V(\lambda) \to 0$ with $d_1(a,b) = ae + b(h-\lambda z)$ and $d_2(b) = b((\lambda+2)z - h, e)$. In particular $V(\lambda)$ is a Koszul $B$-module.*

*Proof.* We look to the map $d_1$ in degree zero: $(d_1)_0(a,b) = ae + b(h-\lambda z) = 0$ with $a, b \in \mathbb{C}$ implies $a = 0$ and $b = 0$, hence $d_1$ is a monomorphism in degree zero.

Let $a$, $b \in B$ be homogeneous elements of degree $\deg ree(a) = degree(b) = m > 0$.

$d_1(a,b) = ae + b(h-\lambda z) = 0$ implies $ae = b(\lambda z - h)$.

The elements $a$, $b$ can be written as:

$a = \sum\limits_{i+j+k+\ell=m} a_{i,j,k,\ell} f^i h^j e^k z^\ell$, $b = \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j e^k z^\ell$.

Then $ae = \sum\limits_{i+j+k+\ell=m} a_{i,j,k,\ell} f^i h^j e^{k+1} z^\ell = \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j e^k z^\ell (\lambda z - h)$.

By induction we prove, $e^k(\lambda z - h) = (2k+\lambda)e^k z - he^k$.

Then $b(\lambda z - h) = \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j ((2k+\lambda)e^k z - he^k) z^\ell =$

$\sum\limits_{i+j+k+\ell=m} (2k+\lambda) b_{i,j,k,\ell} f^i h^j e^k z^{\ell+1} - \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^{j+1} e^k z^\ell$.

Comparing both sides of the equality $ae = b(\lambda z - h)$ we obtain $b_{i,j,0,\ell} = 0$ for $i$, $j$. $\ell$ with $i+j+\ell = m$.

We can cancel $e$ in both sides of the equation to get:

$a = \sum\limits_{i+j+k+\ell=m} a_{i,j,k,\ell} f^i h^j e^k z^\ell =$

$\sum\limits_{i+j+k+\ell=m} (2k+\lambda) b_{i,j,k,\ell} f^i h^j e^{k-1} z^{\ell+1} - \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^{j+1} e^{k-1} z^\ell$.

We prove $he^{k-1} = e^{k-1}h + 2(k-1)e^{k-1}z$.

Substituting in the above equality we have:

$a = \sum\limits_{i+j+k+\ell=m} (2k+\lambda) b_{i,j,k,\ell} f^i h^j e^{k-1} z^{\ell+1} -$

$\sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j e^{k-1} z^\ell\ h - \sum\limits_{i+j+k+\ell=m} 2(k-1) b_{i,j,k,\ell} f^i h^j e^{k-1} z^{\ell+1} =$

$z(\ \sum\limits_{i+j+k+\ell=m} (2+\lambda)\, b_{i,j,k,\ell} f^i h^j e^{k-1} z^\ell) - (\sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j e^{k-1} z^\ell)h$.

Let $b'$ be $b' = \sum\limits_{i+j+k+\ell=m} b_{i,j,k,\ell} f^i h^j e^{k-1} z^\ell = \sum\limits_{i+j+k+\ell=m-1} b_{i,j,k,\ell} f^i h^j e^k z^\ell$.

With this notation, we have proved $a = b'(2+\lambda)z - b'h = b'((\lambda+2)z - h)$ and $b'e = b$, this is $\ker(d_1)_m = b'((\lambda+2)z - h,\ e)$.

Define $d_2(b') = (b'((\lambda+2)z - h),\ b'e)$.

Then the sequence: $0 \to B[-2] \overset{d_2}{\to} B \oplus B[-1] \overset{d_1}{\to} B \to V(\lambda) \to 0$ is exact. ☐

**Corollary 4.** *Let $M(\lambda)$ be a Verma module over the enveloping algebra $U$ of $sl(2,\mathbb{C})$. Then $M(\lambda)$ has global dimension two.*

Since $V(\lambda)$ is a Koszul module, we can apply Koszul duality to obtain a Koszul $B^!$-module $W(\lambda) = \overset{2}{\underset{k=0}{\oplus}} Ext^k_B(V(\lambda), B_0)$. $W(\lambda) = W(\lambda)_0 \oplus W(\lambda)_1 \oplus W(\lambda)_2$ where $W(\lambda)_0 = Hom_{\mathbb{C}}(B/JB, \mathbb{C})$, $W(\lambda)_1 = (Hom_{\mathbb{C}}(B/JB \oplus B/JB, \mathbb{C})) = Hom_{\mathbb{C}}(\mathbb{C}e \oplus \mathbb{C}(h-\lambda z),\mathbb{C})$ $W(\lambda)_2 = Hom_{\mathbb{C}}(B/JB, \mathbb{C})$, with $B/JB \cong \mathbb{C}$.

We want to find the structure of $W(\lambda)$ as $B^!$-module.

In $B^!$ we identify the arrows $e$, $f$, $h$, $z$ with the dual basis $\delta_e$, $\delta_f$, $\delta_h$, $\delta_z$. We start computing the product $e.1$ in $W(\lambda)_0$.

We identify $e$ with the extension $\hat{e} : 0 \to \mathbb{C} \to E \to \mathbb{C} \to 0$.

We have the following commutative diagram:

$$\begin{tikzcd}
0 \arrow[r] & Be + B(h-\lambda z) \arrow[r] \arrow[d, "j"'] & B \arrow[r] \arrow[d, "1"'] & V(\lambda) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & JB \arrow[r] \arrow[d, "p"'] & B \arrow[r] \arrow[dd] & \mathbb{C} \arrow[r] \arrow[dd, "1"'] & 0 \\
 & \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}z \arrow[d, "\delta_e"'] & & & \\
0 \arrow[r] & \mathbb{C} \arrow[r] & E \arrow[r] & \mathbb{C} \arrow[r] & 0
\end{tikzcd}$$

We have a factorization:

$$\begin{tikzcd}
Be + B(h-\lambda z) \arrow[r, "j"] \arrow[dr] & JB \arrow[r, "p"] & \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}z \arrow[dr, "\delta_e"] & \\
 & \mathbb{C}e \oplus \mathbb{C}(h-\lambda z) \arrow[ur, "i"] & & \mathbb{C}
\end{tikzcd}$$

We identify the product $\hat{e}1$ with the map $\delta_e i : \mathbb{C}e \oplus \mathbb{C}(h-\lambda z) \to \mathbb{C}$, where $i$ is the natural inclusion.

The vector space $\mathbb{C}e \oplus \mathbb{C}(h-\lambda z)$ has dual basis $\delta_e$, $\delta_{h-\lambda z}$ and we write $\delta_e i = a\delta_e + b\delta_{h-\lambda z}$, $a$, $b \in \mathbb{C}$.

$\delta_e(h-\lambda z) = a\delta_e(h-\lambda z) + b\delta_{h-\lambda z}(h-\lambda z) = b = 0$ and $\delta_e(e) = a\delta_e(e) + b\delta_{h-\lambda z}(e) = 1 = a$.

We have proved $\hat{e}1 = \delta_e$.

Similarly,$\hat{f}1 = \delta_f i = a\delta_e + b\delta_{h-\lambda z}$ . It follows $\delta_f(e) = 0 = a$, $\delta_f(h - \lambda z = 0 = b$.

Therefore: $\hat{f}1 = 0$.

We compute $\hat{h}1 = \delta_h = a\delta_e + b\delta_{h-\lambda z}$ and $\delta_h(e) = a\delta_e(e) = a = 0$, $\delta_h(h-\lambda z) = 1 = b\delta_{h-\lambda z}(h-\lambda z) = b$.

It follows $\hat{h}1 = \delta_{h-\lambda z}$.

It only remains to compute $\hat{z}1$. As above, $\hat{z}1 = \delta_z = a\delta_e + b\delta_{h-\lambda z}$ and $\delta_z(e) = a\delta_e(e) + b\delta_{h-\lambda z}(e) = a = 0$ and $\delta_z(h-\lambda z)) = b\delta_{h-\lambda z}(h-\lambda z) = b = -\lambda$.

$\hat{z}1 = -\lambda\delta_{h-\lambda z}$.

It follows $(z+\lambda h)1 = -\lambda\delta_{h-\lambda z} + \lambda\delta_{h-\lambda z} = 0$.

Therefore we have the following exact commutative triangle:

$$\begin{tikzcd}
B^! \arrow[rr, "\pi"] \arrow[dr, "q"'] & & W(\lambda) \arrow[r] & 0 \\
 & B^!/B^!f + B^!(z+\lambda h) \arrow[ur, "\varphi"'] & &
\end{tikzcd}$$

the map $\varphi : B^!/B^!f + B^!(z+\lambda h) \to W(\lambda)$ is an epimorphism. Using a dimension argument, we will prove it is an isomorphism.

Since $B^!_0 = \mathbb{C}$, then $(B^!/B^!f + B^!(z+\lambda h))_0 = \mathbb{C}$.

The vector space $B^!_1$ has basis $e$. $f$, $h$, $z$ hence; $e$, $f$, $h$, $(z+\lambda h)$. It follows $(B^!/B^!f + B^!(z+\lambda h))_1$ has basis $\{\overline{e}, \overline{h}\}$.

The vector space $B^!_2$ $\{fe, he, fh, fz, hz, ez\}$,where $fe = -fe$ and $fh = -hf$, hence; $fe, fh \in (B^!f + B^!(z+\lambda h))_2$.

We have $h(z+\lambda h) = hz + \lambda h^2 = hz$ implies $hz \in (B^!f + B^!(z+\lambda h))_2$.

From the equality $fz + zf = 2hf$ it follows $fz = (2h-z)f \in (B^!f + B^!(z+\lambda h))_2$.

Since $e(z+\lambda h) \in (B^!f + B^!(z+\lambda h))_2$ it follows $\overline{ez} = -\lambda\overline{ze}$ in $B^!/B^!f + B^!(z+\lambda h)$.

We have proved $(B^!/B^!f + B^!(z+\lambda h))_2$ is generated by $\overline{he}$.

Now $B^!_3$ is generated by $\{fhe, fez, fhz, hez\}$. It is easy to see that $fhe$, $fez$, $fhz \in B^!f$ and $hez = -ezh$, $-ez(z+\lambda h) = -ez^2 - \lambda ezh = -\lambda ezh$ implies $\overline{hez} = 0$ in $B^!/B^!f + B^!(z+\lambda h)$.

We have proved $(B^!/B^!f + B^!(z+\lambda h))_3 = 0$.

By dimensions, $\varphi$ is an isomorphism.

We have proved:

**Theorem 6.** *For any $\lambda \in \mathbb{C}$ let $W(\lambda)$ be the $B^!$-module corresponding to the homogenized $B$-module $V(\lambda)$ under Koszul duality. Then there is an isomorphism $W(\lambda) \cong B^!/B^!f + B^!(z+\lambda h)$.*

We now have a family $\{W(\lambda)\}_{\lambda\in\mathbb{C}}$ of non isomorphic indecomposable Koszul modules. We know that the graded stable Auslander-Reiten components are of type $ZA_\infty$ [MZ] [Ri]. We will prove below that each $W(\lambda)$ is at the mouth of a regular component, but we need to prove first the following:

**Proposition 9.** *The $B^!$-module $rB^!/socB^!$ is indecomposable.*

*Proof.* We proved above $B^! = \mathbb{C} < e,f,h,z > /I^\perp$ and $I^\perp =< e^2,f^2,h^2,z^2,(e,f),(h,e)$, $(h,f)$, $(h,z)+ef$, $(e,z)-2eh,/f,z)-2hf>$, the algebra $B^!$is graded $B^! = L_0 \oplus L_1 \oplus L_2 \oplus L_3 \oplus L_4$, with $L_0 = \mathbb{C}1$, $L_1 = \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}z$, $L_2 = \mathbb{C}\, fe \oplus \mathbb{C}\, he \oplus \mathbb{C}fh \oplus \mathbb{C}fz \oplus \mathbb{C}hz \oplus \mathbb{C}ez$, $L_3 = \mathbb{C}fhef \oplus \mathbb{C}ez \oplus \mathbb{C}fhz \oplus \mathbb{C}hez$ and $L_4 = \mathbb{C}\, fhz$.

To prove $rB^!/socB^! = L_1 \oplus L_2 \oplus L_3$ is indecomposable we prove that the graded endomorphism ring $Hom_{GrB^!}(\, rB^!/socB^!,\ rB^!/socB^!)$ has dimension one as $\mathbb{C}$-vector space.

Let $E : rB^!/socB^! \to rB^!/socB^!$ be a graded $B^1$-homomorphism.

$E(e) = a_{11}e + a_{12}f + a_{13}h + a_{14}z$

$E(f) = a_{21}e + a_{22}f + a_{23}h + a_{24}z$

$E(h) = a_{31}e + a_{32}f + a_{33}h + a_{34}z$

$E(z) = a_{41}e + a_{42}f + a_{43}h + a_{44}z$

Then $E(e^2) = 0 = eE(e) = a_{12}ef + a_{13}eh + a_{14}ez = -a_{12}fe - a_{13}he + a_{14}z$

Hence; $a_{12} = a_{13} = a_{14} = 0$.

Similarly, $E(f^2) = 0 = fE(f) = a_{21}fe + a_{23}fh + a_{24}fz$ implies $0 = a_{21} = a_{23} = a_{24}$.

$E(h^2) = 0 = hE(h) = a_{31}he + a_{32}hf + a_{34}hz = a_{31}he - a_{32}fh + a_{34}hz$, implies $a_{31} = a_{32} = a_{34} = 0$.

$E(z^2) = 0 = zE(z) = a_{41}ze + a_{42}zf + a_{43}zh = -a_{41}ez - a_{42}fz - a_{43}hz + 2a_{41}eh + 2a_{42}hf - a_{43}ef = -a_{41}ez - a_{42}fz - a_{43}hz - 2a_{41}he - 2a_{42}fh + a_{43}fe$

Implies $a_{41} = a_{42} = a_{43} = 0$.

In the other hand. we have: $E(ef) = -E(fe) = -a_{11}fe$ and $E(ef) = eE(f) = -a_{22}fe$, hence $a_{11} = a_{22}$.
$E(hf) = a_{22}hf$, $E(fh) = a_{33}fh$ , $E(hf) = -E(fh)$ implies $a_{22} = a_{33}$.
$E(zh) = zE(h) = a_{33}zh$ and $E(hz) = hE(z) = a_{44}hz$
$E(zh) = -E(hz) - E(ef) = -a_{44}hz - a_{22}ef = a_{33}zh = a_{33}(-hz - ef)$.
Implies $a_{33} = a_{44}$.
We have proved $a_{11} = a_{22} = a_{33} = a_{44} = a$.
We have now, $E(fhe) = fhE(e) = afhe$.
$E(fez) = feE(z) = afez$, $E(fhz) = fhF(z) = afhz$, $E(ehz) = ehE(z) = aehz$.
Therefore: $E = a1$. □

By $[AR]$ there is an almost split sequence: $0 \to rB^! \to B^! \oplus rB^!/socB^! \to B^!/socB^! \to 0$ and $rB^!$ is at the mouth of a stable component of type $ZA_\infty$.

**Theorem 7.** *For each $\lambda \in \mathbb{C}$ the $B^!$-module $W(\lambda)$ is at the mouth of a regular component.*

*Proof.* We prove first that there are no modules of type $W(\lambda)$ in the preprojective component.

The component of the only indecomposable projective is of the form:



We distinguish five regions in this picture.

Region $A$ consists of all the modules on the two going up diagonals ending either in $rP$ or in $rP/S$ and the irreducible maps corresponding to the arrows. Along these diagonals the arrows represent epimorphisms and all the maps parallel to $rP \to P/S$ are also epimorphisms.

Region $B$ consists of all modules on the going down diagonals containing either $P/S$ or $rP/S$. All the irreducible maps on these diagonals are monomorphism and all maps parallel to $rP/S \to P/S$ are monomorphisms.

Region $C$ consists of all modules below the going down the diagonal containing $rP$ and below the going up diagonal containing $P/S$.In this region the irreducible maps on the going up diagonals are epimorphisms and the irreducible maps in the going down diagonals are monomorphisms.

Region $D$ consists of all modules above the going down diagonal containing $P/S$ and Region $E$ consists of all modules above the going up diagonal containing $rP$.

In both Regions $D$ and $E$ all the irreducible maps on the modules along the going up diagonals are epimorphism and all the irreducible maps of modules along the going down diagonals are monomorphisms.

We will see that a module of the form $W(\lambda)$ can not appear in any of these five regions.

The module $W(\lambda)$ is of the form:
$$\begin{tikzcd}
 & \mathbb{C} & \\
\mathbb{C} \arrow[ur, "1"] & & \mathbb{C} \arrow[ul, "1"'] \\
 & \mathbb{C} \arrow[ul, "1"] \arrow[ur, "1"'] &
\end{tikzcd}$$
and it has as submodules:

$$L = \begin{tikzcd}
 & 0 & \\
\mathbb{C} \arrow[ur] & & \mathbb{C} \arrow[ul] \\
 & \mathbb{C} \arrow[ul, "1"] \arrow[ur, "1"'] &
\end{tikzcd}, \quad L_1 = \begin{tikzcd}
 & 0 & \\
\mathbb{C} \arrow[ur] & & 0 \arrow[ul] \\
 & \mathbb{C} \arrow[ul, "1"] \arrow[ur] &
\end{tikzcd}, \quad L_2 = \begin{tikzcd}
 & 0 & \\
0 \arrow[ur] & & \mathbb{C} \arrow[ul] \\
 & \mathbb{C} \arrow[ul] \arrow[ur, "1"'] &
\end{tikzcd},$$

and
$$L_3 = \begin{tikzcd}
 & 0 & \\
0 \arrow[ur] & & 0 \arrow[ul] \\
 & \mathbb{C} \arrow[ul] \arrow[ur] &
\end{tikzcd}.$$

We have composition series: $W(\lambda) \supset L \supset L_1 \supset L_3$ and $W(\lambda) \supset L \supset L_2 \supset L_3$.

If $W(\lambda)$ appears in Region $D$ it appears at most at tree steps of the mouth, this is $W(\lambda)$ is on a going down diagonal and if it is not at the mouth either $L_3$, $L_1$, $L_2$or $L$ is at the mouth but we can non of the exact sequences: $0 \to L_3 \to L_2 \to S \to 0$, $0 \to L_3 \to L_1 \to S \to 0$, $0 \to L \to W(\lambda) \to S \to 0$ with $S$ simple is almost split.

The reason is that, according to $[AR]$, for an almost split sequence $0 \to \tau(A) \to E \to A \to 0$, $socA \cong top\Omega(A)$ and all modules $L_3$, $L_2$, $L_1$, $L$ have simple socle but $top\Omega S = rB^! / r^2 B^! = \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}z$.

Then if $W(\lambda)$ appears in region $D$ it appears at the mouth.

By the same kind of argument if $W(\lambda)$ appear in region $E$ it appears at the mouth.

But we will see that it is impossible for $W(\lambda)$ to appear at the mouth of the preprojective component.

if it appears in region $D$, then for some integer $k$, $\Omega^{2k}\mathcal{N}^k\ (W(\lambda))$ is isomorphic to $P/S$ where $\mathcal{N}$ denotes the Nakayama equivalence, but this is a contradiction because both the Nakayama equivalence and the syzygy preserve Koszulity and $P/S$ is not Koszul.

If $W(\lambda)$ appears in region $E$, then for some integer $k$ there is an isomorphism $\Omega^{2k}\mathcal{N}^k\ (rP) \cong W(\lambda)$ and this is also impossible since under Koszul duality $\Omega^{2k}\mathcal{N}^k$ $(rP)= F(J^{2k}B)$

and $\dim_{\mathbb{C}} \Omega^{2k}\mathcal{N}^k\ (rP)/r\Omega^{2k}\mathcal{N}^k\ (rP) = \dim_C J2^k/J^{2k+1} = 2k+1 > 1$.

If $W(\lambda)$ appears in region $B$, then it would contain either $rP/S$ or $P/\ S$ which is impossible by dimension.

The module can not appear either in region $A$ because in that case either $rP$ or $rP/S$ would be a quotient of $W(\lambda)$.

Now if $W(\lambda)$ appears in region $C$, then a submodule $L'$ of $W(\lambda)$ appears in the going up diagonal containing $rP/S \to P/S$ and $rP/S$ would be a quotient of $L$, which is impossible by dimensions.

Observe that in a regular component there is only region $D$ and by the above arguments $W(\lambda)$ has to appear at the mouth. $\square$

## 4. Homogenized weight modules

This section will be dedicated to the study of the homogenized weight modules over the homogenized enveloping algebra of $s\ell(2,\,\mathbb{C})$.We will study also a homogenized version of Gelfand's category $\mathcal{O}$ [BGG],[Ma].

Let $M$ be a graded $B$-module. For a given $\lambda \in \mathbb{C}$. we consider the homogenized weight space $V_\lambda$ defined by $V_\lambda = \{m \in M \mid hm = \lambda zm\}$.

We have the following:

**Lemma 2.** *For any graded $B$-module $M$ the $\mathbb{C}$-vector space $\sum\limits_{\lambda\in\mathbb{C}} V_\lambda$ is a $B$-submodule of $M$.*

*Proof.* We claim that given $m \in V_\lambda$ and an integer $k \geq 0$ , $e^k m \in V_{\lambda+2k}$, that $f^k m \in V_{\lambda-2k}$, $h^k \in V_\lambda$ and $z^k m \in V_\lambda$.

We only prove the first claim, the others are similar.

We have proved above $he^k = e^k h + 2ke^k z$.Hence; $(h-\mu z)e^k = e^k(h-(\mu-2k)z)$.

Let $\mu$ be $\mu = \lambda + 2k$. Then $(h-(\lambda+2k)z)e^k m = e^k(h-\lambda z)m = 0$ implies $e^k m \in V_{\lambda+2k}$. □

We are interested in the case the sum is direct.

**Lemma 3.** *Let $M$ be a graded $z$-torsion free $B$-module of the form $M = \sum\limits_{\lambda\in\mathbb{C}} V_\lambda$. Then $M = \bigoplus\limits_{\lambda\in\mathbb{C}} V_\lambda$.*

*Proof.* Assume $0 = m_1 + m_2 + ... + m_k$ with $m_i \in V_{\lambda_i}$ and let $h_i$ be $h_i = (h - \lambda_1 z)(h-\lambda_2 z)...(h-\lambda_{i-1}z)(h-\lambda_{i+1}z)...(h-\lambda_k z)$. Then $h_i m_j = 0$ if $i \neq j$ and $h_i m_i = \prod\limits_{i\neq j}(\lambda_i - \lambda_j)z^{k-1}m_i$.

It follows $h_i 0 = 0 = h_i\ m_1 + h_i m_2 + ... + h_i m_k = \prod\limits_{i\neq j}(\lambda_i-\lambda_j)z^{k-1}m_i$. Since $M$ is $z$-tosion free $m_i = 0$. □

**Definition 2.** *A weight $B$ -module is a graded $B$-module of the form $M = \bigoplus\limits_{\lambda\in\mathbb{C}} V_\lambda$.*

We also need the next lemma.

**Lemma 4.** *Let $M$, $N$ be graded $B$-modules, where $M$ is of the form $M = \sum\limits_{\lambda\in\mathbb{C}} V_\lambda$ and $f : M \to N$ is a surjective homomorphism of $B$-modules. Then $N = \sum\limits_{\lambda\in\mathbb{C}} W_\lambda$ with $W_\lambda = \{n \in N \mid (h-\lambda z)n = 0\}$.*

*Proof.* Let $m \in V_\lambda$. Then $f((h-\lambda z)m) = f(0) = 0 = (h-\lambda z)f(m)$ and $f(V_\lambda) \subset W_\lambda$.

Hence $f(M) = N = \sum\limits_{\lambda\in\mathbb{C}} f(V)_\lambda \subset \sum\limits_{\lambda\in\mathbb{C}} W_\lambda \subset N$ implies $\sum\limits_{\lambda\in\mathbb{C}} W_\lambda = N$. □

**Corollary 5.** *Assume $M = \sum\limits_{\lambda\in\mathbb{C}} V_\lambda$ and let $t_z(M)$ be the $z$-torsion part of $M$. Then $M/\ t_z(M)$ is a weight $B$-module and $M/\ t_z(M) = \bigoplus\limits_{\lambda\in\mathbb{C}} W_\lambda$ with $W_\lambda = f(V_\lambda)$.*

*Proof.* We have a canonical epimorphism $M \to M/\ t_z(M) \to 0$. By the lemma $M/\ t_z(M) = \sum\limits_{\lambda\in\mathbb{C}} W_\lambda$ and by lemma 3, the sum is direct.

We know $f(V_\lambda) \subset W_\lambda$. Let $w \in W_\lambda$. Then $w = f(v_1) + f(v_2) + ...f(v_k)$ with $v_i \in V_{\lambda_i}$ .

Hence $\prod\limits_{i=1}^{k} (h - \lambda_i z)w = \prod\limits_{i=1}^{k} (\lambda - \lambda_i)zw = 0$ implies $\lambda = \lambda_i$ for some $i$.Therefore $0 = f(v_1) + f(v_2) + ...f(v_i) - w + f(v_{i+1}) + ...f(v_k)$ and the fact that the sum is direct implies $f(v_j) = 0$ for $i \neq j$ and $w \in f(V_\lambda)$. ☐

We have the following homogenized version of a result in [Ma].

**Proposition 10.** *Let $M$ be a graded $B$-module of the form $M = \sum\limits_{\lambda\in\mathbb{C}} V_\lambda$, $N$ a graded submodule of $M$ and $W_\lambda = \{n \in N \mid (h - \lambda z)n = 0\}$. Then $N/\sum\limits_{\lambda\in\mathbb{C}} W_\lambda$ is of $z$-torsion.*

*Proof.* Let $n$ be an element in $N$. Then $n = m_1 + m_2 + ...m_k$ with $m_i \in V_{\lambda_i}$

As above $h_i = (h - \lambda_1 z)(h - \lambda_2 z)...(h - \lambda_{i-1} z)(h - \lambda_{i+1} z)...(h - \lambda_k z)$. Then $h_i n = h_i m_i = \prod\limits_{i\neq j} (\lambda_i - \lambda_j)z^{k-1} m_i \in N \cap V_{\lambda_i} = W_{\lambda_i}$.

It follows $z^{k-1} n \in \sum\limits_{\lambda\in\mathbb{C}} W_\lambda$.

We have proved $N/\sum\limits_{\lambda\in\mathbb{C}} W_\lambda$ is of $z$-torsion. ☐

We will consider the subcategory $\mathcal{C}$ of the category $Gr_B$ of graded $B$-modules defined as follows:

A module $M$ is in $\mathcal{C}$ if and only if $M/t_z(M)$ contains a weight submodule $\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ such that the cokernel of the inclusion map: $j : \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \to M/t_z(M)$ is of $z$-torsion.

Informally it means $M$ can be "approximated" by a weight module, or that up to $z$-torsion $M$ is a weight module.

**Definition 3.** *We will call $\mathcal{C}$ the category of generalized weight modules.*

We have the following:

**Proposition 11.** *i) Every submodule of a generalized weight module is a generalized weight module.*

*ii) Each quotient of a generalized weight module is a generalized weight module.*

*iii) A sum of two generalized weight module is a generalized weight module.*

*iii) If $M$, $N$ are two $B$-modules, then $M \otimes_{\mathbb{C}[z]} N$ has structure of $B$-module and if both $M$, $N$ are generalized weight $B$-modules, then $M \otimes_{\mathbb{C}[z]} N$ is a generalized weight $B$-module.*

*Proof.* i) If $N$ is a submodule of $M$, then $t_z(M) \cap N = t_z(N)$.

By hypothesis, there is an exact sequence $0 \to \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \to M/t_z(M) \to X \to 0$, with $X$ of $z$-torsion. Letting $W$ be $W = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \cap (N/t_z(N))$, we have an exact commutative diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & W \arrow[r] \arrow[d] & \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \arrow[r] \arrow[d] & L' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & N/t_z(N) \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] \arrow[d] & L \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & X' \arrow[r] \arrow[d] & X \arrow[r] \arrow[d] & X'' \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

Here $X$ of $z$-torsion implies $X'$ is of $z$-torsion.

Since $W$ is contained in $\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ and it is $z$-torsion free, Lemma 3, and Proposition 10 imply $Y = W/\underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda$ is of $z$-torsion .

We have an exact commutative diagram:

$$\begin{tikzcd}
 & & & 0 \arrow[d] & \\
 & 0 \arrow[d] & 0 \arrow[d] & Y \arrow[d] & \\
0 \arrow[r] & \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda \arrow[r] \arrow[d] & N/t_z(N) \arrow[r] \arrow[d] & Y' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & W \arrow[r] \arrow[d] & N/t_z(N) \arrow[r] \arrow[d] & X' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & Y \arrow[d] & 0 & 0 & \\
 & 0 & & &
\end{tikzcd}$$

where $Y$ and $X'$ are of $z$-torsion. Therefore $Y'$ is of $z$-torsion.

ii) Let $M$ be a generalized weight module and $f : M \to N$ an epimorphism. Then we have the following commutative exact diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & t_z(L) \arrow[r] \arrow[d] & t_z(M) \arrow[r] \arrow[d] & N' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & L \arrow[r] \arrow[d] & M \arrow[r] \arrow[d] & N \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & L/t_z(L) \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] \arrow[d] & N/N' \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

The module $N'$ is of $z$-torsion, hence $N' \subset t_z(N)$ and there is an epimorphism $N/N' \to N/t_z(N) \to 0$. It follows that there is an epimorphism: $M/t_z(M) \overset{\pi}{\to} N/t_z(N) \to 0$.

By hypothesis $M/t_z(M)$ has weight submodule $\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ such the cokernel of the inclusion is of $z$-torsion.

By lemma 4, $\pi(\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda) = \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda$ is a weight submodule of $N/t_z(N)$ and we have an exact commutative diagram:

$$\begin{tikzcd}
0 \arrow[r] & \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] \arrow[d] & X \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda \arrow[r] \arrow[d] & N/t_z(N) \arrow[r] \arrow[d] & X'' \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

and $X$ of $z$-torsion implies $X''$ is of $z$-torsion.

iii) Let $M$, $N$ be $B$-modules in $\mathcal{C}$ . Then there exists exact sequences:

$0 \to \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \to M/t_z(M) \to X \to 0$, $0 \to \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda \to N/t_z(N) \to Y \to 0$, with $X$, $Y$ of $z$-torsion.

Then the sequence:

$0 \to (\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \oplus \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda) \to (M/t_z(M) \oplus N/t_z(N)) \to X \oplus Y \to 0$

Is exact with $X \oplus Y$ of $z$-torsion $(\underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda \oplus \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda) \cong \underset{\lambda\in\mathbb{C}}{\oplus}(V \oplus W)_\lambda$ , $t_z(M) \oplus t_z(N) \cong t_z(M \oplus N)$, where $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$, and $W = \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda$.

Therefore: $M \oplus N$. □

To prove part iv) of the proposition we need some preliminary results.

Since $z$ is in the center $B$ is a $\mathbb{C}[z]$-algebra and the inclusion $\mathbb{C}[z] \to B$ is a morphism of graded $\mathbb{C}$-algebras.

From the existence of a Poincare Birkoff basis of $B$, it follows $B$ is a free $\mathbb{C}[z]$-module.

Let $M$, $N$ be two left $B$-modules. Using the Poincare Birkoff basis we want to define a $B$-module structure on $M \otimes_{\mathbb{C}[z]} N$.

Let $t$ be an element of this basis, this is: $t = e^k f^\ell h^n z^m$, for non negative integers $k$, $\ell$, $n$, $m$.

$\varphi_t : M \times N \to M \otimes_{\mathbb{C}[z]} N$ is the morphism $\varphi_t(m,n) = 1/2(tm \otimes n + m \otimes tn)$.

$\varphi_t$ is bilinear and $\mathbb{C}[z]$-balanced.$\varphi_t(m_1+m_2,n) = 1/2(t(m_1+m_2) \otimes n + (m_1 + m_2) \otimes tn) = 1/2(tm_1 \otimes n + tm_2 \otimes n + (m_1 \otimes n + m_2 \otimes tn) = 1/2(tm_1 \otimes n + m_1 \otimes n)+1/2(tm_2 \otimes n + m_2 \otimes tn) = \varphi_t(m_1,n) + \varphi_t(m_2,n)$.

Similarly, $\varphi_t(m, n_1+n_2) = \varphi_t(m,n_1) + \varphi_t(m,n_2)$ and $\varphi_t(m,zn) = 1/2(tm \otimes zn + m \otimes tzn) = 1/2(tzm \otimes n + zm \otimes tn)$, since $z$ is in the center of $B$.

Hence; $\varphi_t$ induces a morphism $\overline{\varphi_t} : M \otimes_{\mathbb{C}[z]} N \to M \otimes_{\mathbb{C}[z]} N$ given by $\overline{\varphi_t}(m \otimes n) = 1/2(tm \otimes n + m \otimes tn)$. This morphism induces the structure of $B$-module on $M \otimes_{\mathbb{C}[z]} N$.

**Lemma 5.** *Let $M$ be a $z$-torsion free graded $B$-module . Then $M$ is torsion free as $\mathbb{C}[z]$-module.*

*Proof.* Let $m$ be a non zero element of $M$ and $f \in \mathbb{C}[z]$ with $fm = 0$. Since $f$ is non constant $f = (z-\lambda_1)(z-\lambda_2)...(z-\lambda_n)$.

$fm \neq 0$ implies there is an integer $1 < i \leq n$ such that $(z-\lambda_i)...(z-\lambda_n)m = m' \neq 0$ and $(z-\lambda_{i-1})m' = 0$.

Changing $m$ for $m'$ and $\lambda$ for $\lambda_i$, $m \neq 0$ and $(z-\lambda)m = 0$.

Since $M$ is graded $m$ has a decomposition in homogeneous components: $m = m_1 + m_2 + ...m_k$ with $\deg ree(m_i) > \deg ree(m_{i+1})$.

$0 = (z-\lambda)m = zm_1 + zm_2 + ...zm_k - \lambda m_1 - \lambda m_2 ... - \lambda m_k$ and $zm_1 \neq 0$ and of maximal degree since not cancel with any other term in the sum, a contradiction. □

**Corollary 6.** *Let $M$ be graded $z$-torsion free $B$-module. Then $M$ is flat as $\mathbb{C}[z]$-module.*

*Proof.* The module $M$ is a direct limit of finitely generated torsion free modules. In a principal ideal domain a finitely generated torsion free module is free [MB] , hence $M$ is a direct limit of flat modules, then flat. □

**Proposition 12.** *Let $M$ and $N$ be graded $B$-modules, $X$ and $Y$ submodules of $M$ and $N$, respectively, such that $M/X$ and $N/Y$ are $z$-torsion free. Then there are monomorphisms: $X \otimes_{\mathbb{C}[z]} N \to M \otimes_{\mathbb{C}[z]} N$ and $M \otimes_{\mathbb{C}[z]} Y \to M \otimes_{\mathbb{C}[z]} N$ such that $X \otimes_{\mathbb{C}[z]} Y \cong X \otimes_{\mathbb{C}[z]} N \cap M \otimes_{\mathbb{C}[z]} Y$ and $M \otimes_{\mathbb{C}[z]} N/(X \otimes_{\mathbb{C}[z]} N + M \otimes_{\mathbb{C}[z]} Y) \cong (M/X) \otimes_{\mathbb{C}[z]} (N/Y)$.*

*Proof.* Since $M/X$ and $N/Y$ are flat as $\mathbb{C}[z]$-modules, $Tor_1^{\mathbb{C}[z]}(-, N/Y) = 0$ and $Tor_1^{\mathbb{C}[z]}(M/X, -) = 0$, there is an exact commutative diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & X \otimes_{\mathbb{C}[z]} Y \arrow[r] \arrow[d] & X \otimes_{\mathbb{C}[z]} N \arrow[r] \arrow[d] & X \otimes_{\mathbb{C}[z]} N/Y \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & M \otimes_{\mathbb{C}[z]} Y \arrow[r] \arrow[d] & M \otimes_{\mathbb{C}[z]} N \arrow[r] \arrow[d] & M \otimes_{\mathbb{C}[z]} N/Y \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & M/X \otimes_{\mathbb{C}[z]} Y \arrow[d] & M/X \otimes_{\mathbb{C}[z]} N \arrow[d] & M/X \otimes_{\mathbb{C}[z]} N/Y \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

which implies $X \otimes_{\mathbb{C}[z]} Y \cong X \otimes_{\mathbb{C}[z]} N \cap M \otimes_{\mathbb{C}[z]} Y$ and $M \otimes_{\mathbb{C}[z]} N/(X \otimes_{\mathbb{C}[z]} N) \cong M/X \otimes_{\mathbb{C}[z]} N$, $M/X \otimes_{\mathbb{C}[z]} N/(M/X \otimes_{\mathbb{C}[z]} Y) \cong M/X \otimes_{\mathbb{C}[z]} N/Y$.

Hence we have the following chair of isomorphisms:

$$\begin{gathered}
M \otimes_{\mathbb{C}[z]} N/(X \otimes_{\mathbb{C}[z]} N + M \otimes_{\mathbb{C}[z]} Y) \cong \\
\left(M \otimes_{\mathbb{C}[z]} N/X \otimes_{\mathbb{C}[z]} N\right)/((X \otimes_{\mathbb{C}[z]} N + M \otimes_{\mathbb{C}[z]} Y)/(X \otimes_{\mathbb{C}[z]} N) \cong \\
(M/X \otimes_{\mathbb{C}[z]} N)/((M \otimes_{\mathbb{C}[z]} Y)/X \otimes_{\mathbb{C}[z]} NY)) \cong \\
M/X \otimes_{\mathbb{C}[z]} N)/M/X \otimes_{\mathbb{C}[z]} N) \cong M/X \otimes_{\mathbb{C}[z]} N/Y
\end{gathered}$$

□

We will also need the following:

**Proposition 13.** *Let $M$, $N$ be graded $z$-torsion free $B$-modules. Then $M \otimes_{\mathbb{C}[z]} N$ is $z$-torsion free.*

*Proof.* Let $\sum\limits_{i=1}^{k} m_i \otimes n_i$ be a non zero element of $M \otimes_{\mathbb{C}[z]} N$ and $\ell > 0$ such that $z^\ell \sum\limits_{i=1}^{k} m_i \otimes n_i = 0$. We may assume $m_i$, $n_i$ are homogeneous.

$\sum\limits^{k} \mathbb{C}[z] n_i$ is a finitely generated $z$-torsion free submodule of $N$, hence it is a free $\mathbb{C}[z]$-module.

Let $\varphi : \bigoplus\limits_{j=1}^{t} \mathbb{C}[z] \to \sum\limits^{k} \mathbb{C}[z] n_i$ be an isomorphism and $e_j = (0, 0...0, 1, 0...0)$ the canonical basis of $\bigoplus\limits_{j=1}^{t} \mathbb{C}[z]$.

The morphism $\varphi$ induces an isomorphism: $1\otimes\varphi : M\otimes_{\mathbb{C}[z]} \bigoplus_{j=1}^{t}\mathbb{C}[z] \to M\otimes_{\mathbb{C}[z]} \sum_{i=1}^{k}\mathbb{C}[z]n_i$.

Since $M$ is a flat $\mathbb{C}[z]$-module, the inclusion $j : \sum_{i=1}^{k}\mathbb{C}[z]n_i \to N$ induces a monomorphism $1\otimes j : M\otimes_{\mathbb{C}[z]} \sum_{i=1}^{k}\mathbb{C}[z]n_i \to M\otimes_{\mathbb{C}[z]} N$.

We have $\varphi^{-1}(n_j) = \sum_{i=1}^{t}c_{ij}e_i$ with $c_{ij}\in\mathbb{C}[z]$.

Then $(1\otimes j)(1\otimes\varphi)(\sum_{i=1}^{k}m_i\otimes(\sum_{i=1}^{t}c_{ij}e_i) = \sum_{i=1}^{k}m_i\otimes n_i$.

$\sum_{i=1}^{k}m_i\otimes(\sum_{i=1}^{t}c_{ij}e_i) = \sum_{i=1}^{k}\ \sum_{i=1}^{t}m_ic_{ij}\otimes e_i$.

Then $z^\ell\sum_{i=1}^{k}m_i\otimes n_i = 0 = z^\ell(1\otimes j)(1\otimes\varphi)(\sum_{i=1}^{k}m_i\otimes(\sum_{i=1}^{t}c_{ij}e_i)) = \sum_{i=1}^{k}\ z^\ell(\sum_{i=1}^{t}m_ic_{ij})\otimes e_i$ implies $z^\ell(\sum_{i=1}^{t}m_ic_{ij}) = 0$ and $M$ $z$-torsion free implies $\sum_{i=1}^{t}m_ic_{ij} = 0$.

Therefore: $\sum_{i=1}^{k}m_i\otimes n_i = (1\otimes j)(1\otimes\varphi)(\sum_{i=1}^{k}\ \sum_{i=1}^{t}m_ic_{ij}\otimes e_i) = 0$.

We have proved $M\otimes_{\mathbb{C}[z]} N$ is $z$-torsion free. □

Applying the previous results we get:

**Lemma 6.** *Let $M$ and $N$ be graded $B$-modules and denote by $t_z(-)$ the radical corresponding to $z$- torsion. Then $t_z(M\otimes_{\mathbb{C}[z]} N) = t_z(M)\otimes_{\mathbb{C}[z]} N + M\otimes_{\mathbb{C}[z]} t_z(N)$.*

*Proof.* Since $N/t_z(N)$ and $M/t_z(M)$ are flat $\mathbb{C}[z]$-modules, the exact sequences:

$$0\to\ t_z(M)\ \to\ M\ \to\ M/t_z(M)\ \to 0\ ,$$

$$0\to\ t_z(N)\ \to\ N\ \to\ N/t_z(N)\ \to 0$$

Induce by tensoring exact sequences:

$$0\to\ t_z(M)\otimes_{\mathbb{C}[z]} N\ \to\ M\otimes_{\mathbb{C}[z]} N\ \to\ M/t_z(M)\otimes_{\mathbb{C}[z]} N\ \to 0\ ,$$

$$0\to\ M\otimes_{\mathbb{C}[z]} t_z(N)\ \to\ M\otimes_{\mathbb{C}[z]} N\ \to\ M\otimes_{\mathbb{C}[z]} N/t_z(N)\ \to 0$$

Since $t_z(M)\otimes_{\mathbb{C}[z]} N$, $M\otimes_{\mathbb{C}[z]} t_z(N)$ are $z$-torsion modules, $t_z(M)\otimes_{\mathbb{C}[z]} N + M\otimes_{\mathbb{C}[z]} t_z(N)$ is a $z$-torsion submodule of $M\otimes_{\mathbb{C}[z]} N$, hence it is obtained in $t_z(M\otimes_{\mathbb{C}[z]} N)$.

In the other hand, $M\otimes_{\mathbb{C}[z]} N/(\ t_z(M)\otimes_{\mathbb{C}[z]} N + M\otimes_{\mathbb{C}[z]} t_z(N)) \cong M/t_z(M)\otimes_{\mathbb{C}[z]} N/t_z(N)$.

By Proposition 13, $M/t_z(M)\otimes_{\mathbb{C}[z]} N/t_z(N)$ is $z$-torsion free. It follows, $t_z(M\otimes_{\mathbb{C}[z]} N) = t_z(M)\otimes_{\mathbb{C}[z]} N + M\otimes_{\mathbb{C}[z]} t_z(N)$. □

After these results, we can finally prove:

**Claim 1.** *Let $M$ and $N$ be generalized weight $B$-modules. Then $M\otimes_{\mathbb{C}[z]} N$ is a generalized weight $B$-modules.*

*Proof.* By hypothesis, we have exact sequences:

$$0 \to \bigoplus_{\lambda\in\mathbb{C}} V_\lambda \to M/t_z(M) \to Z \to 0$$

$$0 \to \bigoplus_{\lambda\in\mathbb{C}} W_\lambda \to N/t_z(N) \to R \to 0$$

with $Z$, $R$, $z$-torsion modules.

Since $\bigoplus_{\lambda\in\mathbb{C}} V_\lambda$, $\bigoplus_{\lambda\in\mathbb{C}} W_\lambda$, $M/t_z(M)$, $N/t_z(N)$ are flat $\mathbb{C}[z]$-modules, we have the following commutative exact diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & \bigoplus_{\lambda\in\mathbb{C}} \mathrm{V}_\lambda \otimes_{\mathbb{C}[z]} \bigoplus_{\lambda\in\mathbb{C}} \mathrm{W}_\lambda \arrow[r] \arrow[d] & \mathrm{M}/\mathrm{t}_z(\mathrm{M}) \otimes_{\mathbb{C}[z]} \bigoplus_{\lambda\in\mathbb{C}} \mathrm{W}_\lambda \arrow[r] \arrow[d] & \mathrm{Z} \otimes_{\mathbb{C}[z]} \bigoplus_{\lambda\in\mathbb{C}} \mathrm{W}_\lambda \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \bigoplus_{\lambda\in\mathbb{C}} \mathrm{V}_\lambda \otimes_{\mathbb{C}[z]} \bigoplus_{\lambda\in\mathbb{C}} \mathrm{W}_\lambda \arrow[r] \arrow[d] & \mathrm{M}/\mathrm{t}_z(\mathrm{M}) \otimes_{\mathbb{C}[z]} \mathrm{N}/\mathrm{t}_z(\mathrm{N}) \arrow[r] \arrow[d] & L \arrow[r] \arrow[d] & 0 \\
 & 0 \arrow[r] & \mathrm{M}/\mathrm{t}_z(\mathrm{M}) \otimes_{\mathbb{C}[z]} \mathrm{R} \arrow[r] \arrow[d] & \mathrm{M}/\mathrm{t}_z(\mathrm{M}) \otimes_{\mathbb{C}[z]} \mathrm{R} \arrow[r] \arrow[d] & 0 \\
 & & 0 & 0 &
\end{tikzcd}$$

Since $Z \otimes_{\mathbb{C}[z]} \bigoplus_{\lambda\in\mathbb{C}} W_\lambda$ and $M/t_z(M) \otimes_{\mathbb{C}[z]} R$ are of $z$-torsion, $L$ is of $z$-torsion.

Letting $V$ and $W$ be the weight spaces $V = \bigoplus_{\lambda\in\mathbb{C}} V_\lambda$ and $W = \bigoplus_{\lambda\in\mathbb{C}} W_\lambda$ we only need to prove that $V \otimes_{\mathbb{C}[z]} W = \bigoplus_{\lambda\in\mathbb{C}} (V \otimes_{\mathbb{C}[z]} W)_\lambda$.

We will actually prove $(V \otimes_{\mathbb{C}[z]} W)_\lambda = \bigoplus_{\lambda=\mu+\sigma} V_\mu \otimes_{\mathbb{C}[z]} W_\sigma$.

We have a decomposition as $\mathbb{C}[z]$-module: $V \otimes_{\mathbb{C}[z]} W = \bigoplus_{\mu,\sigma\in\mathbb{C}} V_\mu \otimes_{\mathbb{C}[z]} W_\sigma$.

Let $v$ be an element of $(V \otimes_{\mathbb{C}[z]} W)_\lambda$.Then $v = v_1 + v_2 + ...v_k$ with $v_i \in V_{\mu_i} \otimes_{\mathbb{C}[z]} W_{\sigma_i}$, this is: $v_i = \sum_{j=1}^{t_i} v_j^{(i)} \otimes w_j^{(i)}$ , $v_j^{(i)} \in V_{\mu_i}$ and $w_j^{(i)} \in W_{\sigma_i}$.

Using the structure of $V \otimes_{\mathbb{C}[z]} W$ as $B$-module, $h\ v_i = 1/2(\sum_{j=1}^{t_i} hv_j^{(i)} \otimes w_j^{(i)} + \sum_{j=1}^{t_i} v_j^{(i)} \otimes hw_j^{(i)}) = z/2(\sum_{j=1}^{t_i} \mu_i v_j^{(i)} \otimes w_j^{(i)} + \sum_{j=1}^{t_i} v_j^{(i)} \otimes \sigma_j w_j^{(i)}) = (\mu_i + \sigma_j) z \sum_{j=1}^{t_i} v_j^{(i)} \otimes w_j^{(i)} = (\mu_i + \sigma_j) z v_i$.

Hence $hv = hv_1 + hv_2 + ...hv_k = z((\mu_1 + \sigma_1)v_1 + (\mu_2 + \sigma_2)v_2 + ...(\mu_k + \sigma_k)v_k) = \lambda z v = z(\lambda v_1 + \lambda v_2 + ...\lambda v_k$.

Therefore $z(\lambda - (\mu_i + \sigma_i)v_i = 0$ , $V \otimes_{\mathbb{C}[z]} W$ is $z$-torsion free. It follows $\lambda = \mu_i + \sigma_i$.

We have proved $(V \otimes_{\mathbb{C}[z]} W)_\lambda \subset \bigoplus_{\lambda=\mu+\sigma} V_\mu \otimes_{\mathbb{C}[z]} W_\sigma$.

Now let $v \in V_\mu$ and $w \in W_\sigma$ with $\lambda = \mu + \sigma$.

Then $h(v \otimes w) = 1/2(hv \otimes w + v \otimes hw) = (\mu + \sigma) z (v \otimes w) = \lambda z (v \otimes w)$.

Then $v \otimes w \in (V \otimes_{\mathbb{C}[z]} W)_\lambda$.

We have proved $(V \otimes_{\mathbb{C}[z]} W)_\lambda = \bigoplus_{\lambda=\mu+\sigma} V_\mu \otimes_{\mathbb{C}[z]} W_\sigma$. □

**Definition 4.** *The homogenized category $\mathcal{O}_B$ is the subcategory of the category of generalized weight $B$-modules $\mathcal{C}$ satisfying the following two conditions:*

*i) The module $M$ in $\mathcal{O}_B$ is finitely generated.*

*ii) For all $v \in M$, $dim \mathbb{C}[z]v < \infty$.*

We will see below that the homogenized Verma modules form a subcategory of $\mathcal{O}_B$.

**Theorem 8.** *The category* $\mathcal{O}_B$ *is closed under submodules, quotients and finite direct sums. In particular is an abelian Krull-Schmidt category.*

**Remark 1.** *Since* $B$ *is a Koszul algebra of finite global dimension, for a given finitely generated* $B$*-module* $M$ *there is a truncation* $M_{\geq k}[k]$ *which is Koszul in particular if* $M$ *is in* $\mathcal{O}_B$ *the Koszul module* $M_{\geq k}[k]$ *is in* $\mathcal{O}_B$*.*

*Moreover, there is a commutative exact diagram:*

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & t_z(M_{\geq k}) \arrow[r] \arrow[d] & t_z(M) \arrow[r] \arrow[d] & X \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & M_{\geq k} \arrow[r] \arrow[d] & M \arrow[r] \arrow[d] & M/M_{\geq k} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & M_{\geq k}/t_z(M_{\geq k}) \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] \arrow[d] & Y \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

*with both* $X,\ Y$ *finite dimensional* $\mathbb{C}$*-vector spaces.*

**Proposition 14.** *Every homogenized Verma module* $V(\lambda)$ *is in* $\mathcal{O}_B$*.*

*Proof.* We proved above that the graded $B$-module $V(\lambda)$ is defined as $V(\lambda) = B/(Be+B(h-\lambda z))$ and that as $C[z]$-module it decomposes $V(\lambda) = \underset{k\geq 0}{\oplus}(\underset{i+j=k}{\oplus}\mathbb{C}f^iz^j)$.

An element $b$ of $V(\lambda)$ has a decomposition in homogeneous components: $b = b_{i_1} + b_{i_2} + ...b_{i_k}$ with $b_{i_j}$ homogeneous of degree $m_j$.

We have: $(h-\mu z)b = 0 = (h-\mu z)b_{i_1} + (h-\mu z)b_{i_2} + ...(h-\mu z)b_{i_k}$ if and only if for all $j$, $(h-\mu z)b_{ij} = 0$.

We may assume $b$ is homogeneous of degree $k$.Then $b = \sum\limits_{i+j=k} c_{ij}f^iz^j$ with $c_{ij} \in \mathbb{C}$.

$(h-\mu z)f^i = hf^i - \mu zf^i = f^ih - 2if^iz - \mu zf^i = f^i(h-(2i+\mu)z)$.

It follows $(h-\mu z)f^i = 0$ in $V(\lambda)$ if and only if $\lambda = 2i+\mu$.

Therefore: $(h-(\lambda-2i)z)f^iz^j = 0$ for all $i$,$j$, $i+j=k$.

Since $V(\lambda)$ is $z$-torsion free it follows $V(\lambda) = \underset{i\geq 0}{\oplus} V_{\lambda-2i}$ is a weight module.

By induction we have the following equality: $ef^n = f^ne + nf^{n-1}z(h-\lambda z) + n(\lambda-(n-1))f^{n-1}z^2$.

Hence; $ef^n + (Be+B(h-\lambda z)) = n(\lambda-(n-1))f^{n-1}z^2 + (Be+B(h-\lambda z))$.

By induction we have for $k \leq n$, $e^kf^n + (Be+B(h-\lambda z)) = n(n-1)...(n-(k-1))(\lambda-(n-1))((\lambda-(n-2))...(\lambda-(n-k))f^{n-k}z^{2k} + (Be+B(h-\lambda z))$.

In particular for $k=n$, $e^nf^n + (Be+B(h-\lambda z)) = n(n-1)(\lambda-(n-1))((\lambda-(n-2))...(\lambda-1)\lambda z^{2n} + (Be+B(h-\lambda z))$.

Therefore: $e^{n+1}f^n = 0$ in $V(\lambda)$.

It follows that for any element $f^iz^j$ of $V(\lambda)$, $e^{i+1}f^iz^j = 0$.

From this it follows $V(\lambda)$ is in $\mathcal{O}_B$. □

4.1. **Weight Modules.** In this subsection we continue the study of weight $B$-modules, and concentrate in those that are $z$-torsion free and belong to the category $\mathcal{O}_B$. For the full subcategory of $\mathcal{O}_B$ containing such modules we define a duality.

Let $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ be a graded weight $B$-module. Then $V$ decomposes in homogeneous components as $V = \underset{i\in\mathbb{Z}}{\oplus} V_i$. We compare both decompositions. Let $v \in V_\lambda$ and $v = v_{i_1} + v_{i_2} + ...v_{i_k}$ be a decomposition in homogeneous components $v_{ij} \in V_{i_j}$ with $i_1 < i_2 < ... < i_k$.

$(h-\lambda z)v = 0 = (h-\lambda z)v_{i_1} + (h-\lambda z)v_{i_2} + ...(h-\lambda z)v_{i_k}$ and $(h-\lambda z)v_{ij} \in V_{i_j+1}$ implies $(h-\lambda z)v_{ij} = 0$ for $1 \leq j \leq k$. and $v_{ij} \in V_{ij} \cap V_\lambda$.

We have $\sum\limits_i V_\lambda \cap V_i \subset V_\lambda \subset \sum\limits_i V_\lambda \cap V_i$, therefore: $\underset{i\in\mathbb{Z}}{\oplus} V_\lambda \cap V_i = V_\lambda$.

For the other decomposition $\sum\limits_i V_\lambda \cap V_i \subset V_i$.

Let $v$ be an element of $V_\ell$ and take its decomposition $v = v_1 + v_2 + ...v_k$ with $v_i \in V_{\lambda_i} = \underset{j\in\mathbb{Z}}{\oplus} V_{\lambda_i} \cap V_j$.

Then each $v_i = \sum\limits_{j=1}^{t_i} w_j^{(i)}$ and $w_j^{(i)} \in V_{\lambda_i} \cap V_j$ and degree$w_j^{(i)} = j$.

We write $v$ as $v = \sum\limits_{i=1}^{k}\sum\limits_{j=1}^{t_i} w_j^{(i)} = \sum\limits_{j=1}^{t_i}\sum\limits_{i=1}^{k} w_j^{(i)}$. It follows $\sum\limits_{i=1}^{k} w_j^{(i)}$ is homogeneous of degree $j$ .

Since $v$ is homogeneous of degree $\ell$ the element $\sum\limits_{i=1}^{k} w_j^{(i)} = 0$ for $j \neq \ell$. and $v = \sum\limits_{i=1}^{k} w_\ell^{(i)}$ with $w_\ell^{(i)} \in V_\ell \cap V_{\lambda_i}$.

We proved $V_\ell \subset \sum\limits_{\ell\in\mathbb{C}} V_\lambda \cap V_\ell$.It follows $V_\ell = \underset{\ell\in\mathbb{C}}{\oplus} V_\lambda \cap V_\ell$.

Then $V = \underset{\lambda\in\mathbb{C}}{\oplus}\ \underset{i\geq 0}{\oplus} (V_\lambda \cap V_i)$.

As a consequence we have: $V_{\geq k} = \underset{i\geq k}{\oplus} V_i = \underset{i\geq k}{\oplus}(\underset{\lambda\in\mathbb{C}}{\oplus}(V_\lambda \cap V_i)) = \underset{\lambda\in\mathbb{C}}{\oplus}(\underset{i\geq k}{\oplus}(V_\lambda \cap V_i))$ and $(V_{\geq k})_\lambda = \underset{i\geq k}{\oplus}(V_\lambda \cap V_i)$.

It follows $V_{\geq k} = \underset{\lambda\in\mathbb{C}}{\oplus}(V_{\geq k})_\lambda$, this is: any truncation of a weight module is a weight module.

We have actually proved:

**Theorem 9.** *Let* $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ *be a graded weight* $B$*-module. Then there is a truncation* $V_{\geq k}$ *which is Koszul and* $V_{\geq k}$ *is a weight module.*

**Corollary 7.** *Let* $M$ *be a finitely generated generalized weight* $B$*-module. Then there exists a weight submodule* $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ *of* $M/t_z(M)$ *such that for some integer* $k$*,* $V[k]$ *is Koszul and* $(M/t_z(M))/V$ *is of* $z$*-torsion.*

*Proof.* By definition, there is an exact sequence: $0 \to \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda \to M/t_z(M) \to R \to 0$ with $R$ of $z$-torsion. Since $M/t_z(M)$ is finitely generated and $B$-noetherian, $W = \underset{\lambda\in\mathbb{C}}{\oplus} W_\lambda$ is finitely generated and there is a truncation $V = W_{\geq k}$ with $V[k]$ Koszul and $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ a weight module,

We have an exact commutative diagram;

$$\begin{tikzcd}
 & & & 0 \arrow[d] & \\
 & 0 \arrow[d] & & L \arrow[d] & \\
0 \arrow[r] & V \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] \arrow[d, "1"'] & X \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & W \arrow[r] \arrow[d] & M/t_z(M) \arrow[r] & R \arrow[r] \arrow[d] & 0 \\
 & L \arrow[d] & & 0 & \\
 & 0 & & &
\end{tikzcd}$$

with $L$ a finite dimensional $\mathbb{C}$-vector space and $R$ of $z$-torsion.

Therefore: $X$ is of $z$-torsion. □

**Theorem 10.** *Let $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ be a $z$-torsion free weight module and assume $V$ is in the category $\mathcal{O}_B$. Then for each $\lambda \in \mathbb{C}$, $V_\lambda$ is a finitely generated $\mathbb{C}[z]$-module, in particular each $V_\lambda$ is a free $\mathbb{C}[z]$-module of finite rank.*

*Proof.* We have proved the following equalities: $he^n = e^n h + 2ne^n z$ and $hf^n = f^n h - 2nf^n z$.

By hypothesis $V$ is a finitely generated $B$-module and for each $v \in V$ there is a positive integer $\ell$ such that $e^\ell v = 0$.

We have decompositions of $V$: $V = \underset{i\geq k_0}{\oplus} V_i$ in homogeneous components and $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ in weight spaces and $V = \underset{\lambda\in\mathbb{C}}{\oplus}\,\underset{i\geq k_0}{\oplus} V_\lambda \cap V_i$.

Hence; we can assume $V$ is generated by homogeneous elements $v_{i_1}$, $v_{i_2}$,... $v_{i_k}$ of degree $v_{i_j} = i_j$ and $(h - \lambda_j z)v_{i_j} = 0$.

For each $j$ there is an $\ell_j$ such that $e^{\ell_j} v_{i_j} = 0$, then for $\ell = \max\{\ell_j\}$, $e^\ell v_{i_j} = 0$ for all $j$.

Let $v$ be an element of $V_\lambda$.Then $v = w_{k_0} + w_{k_0+1} + ...w_{k_0+n}$ is a decomposition in homogeneous components $w_{k_0+j} \in V_{k_0+j} \cap V_\lambda$.

We write $w_{k_0+t} = \sum\limits_{j=1}^{k} b_{tj} v_{i_j}$ with $b_{tj} = \sum\limits_{s+r+n+m=t-i_j} c^{(j)}_{s,r,n,m} f^s e^r h^n z^m$ and $c^{(j)}_{s,r,n,m} \in \mathbb{C}$. Then $w_{k_0+t} = \sum\limits_{j=1}^{k} \sum\limits_{s+r+n+m=t-i_j} c^{(j)}_{s,r,n,m} f^s e^r h^n z^m v_{i_j}$.

Hence each $w_{k_0+t}$ is a linear combination of the vectors$\{ f^s e^r h^n z^m v_{i_j}\}$ and changing for a smaller set if necessary we my assume they are linearly independent and $w_{k_0+t}$ is still a linear combination of such vectors.

We have $h^n v_{i_j} = \lambda_j^n z^n v_{i_j}$, hence $w_{k_0+t} = \sum\limits_{j=1}^{k} \sum\limits_{s+r+n+m=t-i_j} \lambda_j^n c^{(j)}_{s,r,n,m} f^s e^r z^{m+n} v_{i_j}$.

From the equalities: $(h - \lambda z)f^s e^r = f^s e^r (h - (\lambda + 2(s-r))z)$ and $(h - (\lambda + 2(s-r))z)v_{i_j} = (\lambda_j - (\lambda + 2(s-r)))zv_{i_j}$ it follows:

$$0 = (h - \lambda z\ w_{k_0+t} = z \sum_{j=1}^{k} \sum_{s+r+n+m=t-i_j} (\lambda_j - (\lambda + 2(s-r)))\lambda_j^n c^{(j)}_{s,r,n,m} f^s e^r z^{m+n} v_{i_j}.$$

Since $V$ is $z$-torsion free $\sum\limits_{j=1}^{k}\sum\limits_{s+r+n+m=t-i_j}(\lambda_j-(\lambda+2(s-r)))\lambda_j^n c_{s,r,n,m}^{(j)} f^s e^r z^{m+n} v_{i_j} = 0$ . By the hypothesis that they are linearly independent $\lambda_j = \lambda + 2(s-r)$ and $v_{i_j} \in V_{\lambda+2(s-r)}$. Therefore: $f^s e^r v_{i_j} \in V_\lambda$ with $0 \leq r \leq \ell - 1$, $s \geq 0$.

If $f^{s'} e^{r'} v_{i_j}$ is another element of $V_\lambda$, then $\lambda = \lambda_j + 2(s'-r')$. It follows $s' = s + r' - r$ with $0 \leq r' \leq \ell - 1$ and $f^{s'} e^{r'} v_{i_j} = f^{s+r'-r} e^{r'} v_{i_j}$. For any pair of fixed elements ( $s, r$) $0 \leq r \leq \ell - 1$, $s \geq 0$ , there are only a finite number of elements of the form $f^{s+r'-r} e^{r'} v_{i_j} = f^{s'} e^{r'} v_{i_j}$ in $V_\lambda$.

This implies $w_{k_0+t}$ is a polynomial combination $w_{k_0+t} = \sum q(z)_{s,r,i_j} f^s e^r v_{i_j}$ with $q(z)_{s,r,i_j} \in \mathbb{C}[z]$.

It follows $V_\lambda$ is a finitely generated $\mathbb{C}[z]$-module and $V$ torsion free implies $V_\lambda$ is a free $\mathbb{C}[z]$-module of finite rank. □

Let $M$ be an object in $\mathcal{O}_B$. Then there is a weight $B$-module $V$ and an exact sequence:

$$0 \to \quad V \quad \overset{i}{\to} \quad M/t_z(M) \quad \overset{p}{\to} \quad R \quad \to 0$$

with $R$ of $z$-torsion, which induces an exact sequence:

$$0 \to \quad V_\lambda \quad \overset{i}{\to} \quad (M/t_z(M))_\lambda \quad \overset{p}{\to} \quad p((M/t_z(M))_\lambda) \quad \to 0$$

with $p((M/t_z(M))_\lambda)$ a $\mathbb{C}[z]$ -submodule of $R$, hence of $z$-torsion. There is an integer $k \geq 0$ such that $z^k p((M/t_z(M))_\lambda) = 0$. It follows $z^k(M/t_z(M))_\lambda) \subset V_\lambda$

But $(M/t_z(M))_\lambda$ $z$-torsion free, implies the map given by multiplication $z^k : (M/t_z(M))_\lambda \to z^k((M/t_z(M))_\lambda$ is an isomorphism and there is a monomorphism $z^k : (M/t_z(M))_\lambda \to V_\lambda$ hence, $(M/t_z(M))_\lambda$ is a finitely generated torsion free $\mathbb{C}[z]$-module. We have proved the first part of the following:

**Proposition 15.** *Let $M$ be an object in $\mathcal{O}_B$. Then:*

*i) For any $\lambda \in \mathbb{C}$ the $\mathbb{C}[z]$-module $(M/t_z(M))_\lambda$ is either zero or it is a free $\mathbb{C}[z]$-module of finite rank.*

*ii) There is a $\mathbb{C}[z]$-module decomposition $M_\lambda = t_z(M_\lambda) \oplus M_\lambda/t_z(M_\lambda)$ where $M_\lambda/t_z(M_\lambda)$ is either zero or it is a free $\mathbb{C}[z]$-module of finite rank.*

*Proof.* ii) We have the following commutative exact diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & t_z(M)_\lambda \arrow[r] \arrow[d] & M_\lambda \arrow[r] \arrow[d] & p(M_\lambda) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & t_z(M) \arrow[r, "i"] & M \arrow[r, "p"] & M/t_z(M) \arrow[r] & 0
\end{tikzcd}$$

where $p(M_\lambda) \subset (M/t_z(M))_\lambda$. It is clear that $t_z(M_\lambda) = M_\lambda \cap t_z(M) = t_z(M)_\lambda$. Therefore: $p(M_\lambda) = M_\lambda/t_z(M_\lambda)$

By the first part, $(M/t_z(M))_\lambda$ is either zero or a free $\mathbb{C}[z]$-module of finite rank. In the first case $p(M_\lambda) = M_\lambda/t_z(M_\lambda) = 0$ and the decomposition holds.

In the second case $p(M_\lambda) = M_\lambda/t_z(M_\lambda)$ is a free $\mathbb{C}[z]$-module of finite rank and we have again the decomposition $M_\lambda = t_z(M_\lambda) \oplus M_\lambda/t_z(M_\lambda)$. □

Assume now $M \in \mathcal{O}_B$ is $z$-torsion free. By hypothesis there is a weight submodule $V$ and an exact sequence: $0 \to V \to M \to R \to 0$ with $V = \underset{\lambda\in\mathbb{C}}{\oplus} V_\lambda$ and $R$ of

$z$ -torsion.. Since $V_\lambda \subset M_\lambda$ implies $\sum\limits_{\lambda\in\mathbb{C}} V_\lambda \subset \sum\limits_{\lambda\in\mathbb{C}} M_\lambda$ and $M$ $z$-torsion free implies $\sum\limits_{\lambda\in\mathbb{C}} M_\lambda = (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda$.

There is an epimorphism: $M/V \to M/(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda) \to 0$ and $M/(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)$ is also of $z$-torsion. Hence $\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda$is another weight module approximating $M$ and we may assume $V = \underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda$ where each $M_\lambda$ is either zero or a free $\mathbb{C}[z]$-module of finite rank.

We that for $m \in M_\lambda$, $e^k m \in M_{\lambda+2k}$, $f^k \in M_{\lambda-2k}$, $h^k \in M_\lambda$, $z^k \in M_\lambda$. Let $M_\lambda^*$ be $M_\lambda^* = Hom_{\mathbb{C}[z]}(M_\lambda, \mathbb{C}[z])$ and define $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^* = \underset{\lambda\in\mathbb{C}}{\oplus} Hom_{\mathbb{C}[z]}(M_\lambda, \mathbb{C}[z]) = \underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda^*$ .

Given a map $\varphi = (0, 0 ... \varphi_{\lambda_1},\ \varphi_{\lambda_2},\ ...\varphi_{\lambda_r} 0,\ 0, ....\ ) \in (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$.

We define $e^k\varphi = (0, 0, ..e^k\varphi_{\lambda_1},\ e^k\varphi_{\lambda_2},\ ...e^k\varphi_{\lambda_r} 0,\ 0, ....\ )$, where $e^k\varphi_{\lambda_i} : M_{\lambda_i - 2k} \to \mathbb{C}[z]$ given by $e^k\varphi_{\lambda_i}(m) = \varphi_{\lambda_i}(e^k m)$.

Similarly $f^k\varphi = (0, 0, ..f^k\varphi_{\lambda_1},\ f^k\varphi_{\lambda_2},\ ...f^k\varphi_{\lambda_r} 0,\ 0, ....\ )$, with $f^k\varphi_{\lambda_i} : M_{\lambda_i + 2k} \to \mathbb{C}[z]$ given by $f^k\varphi_{\lambda_i}(m) = \varphi_{\lambda_i}(f^k m)$.

$h^k\varphi = (0, 0, ..h^k\varphi_{\lambda_1},\ h^k\varphi_{\lambda_2},\ ...h^k\varphi_{\lambda_r} 0,\ 0, ....\ )$, with $h^k\varphi_{\lambda_i} : M_{\lambda_i} \to \mathbb{C}[z]$ given by $h^k\varphi_{\lambda_i}(m) = \varphi_{\lambda_i}(h^k m)$.

$z^k\varphi = (0, 0, ..z^k\varphi_{\lambda_1},\ z^k\varphi_{\lambda_2},\ ...z^k\varphi_{\lambda_r} 0,\ 0, ....\ )$, with $z^k\varphi_{\lambda_i} : M_{\lambda_i} \to \mathbb{C}[z]$ given by $z^k\varphi_{\lambda_i}(m) = \varphi_{\lambda_i}(z^k m)$.

With these operations $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$ becomes a $B$-module.

We claim $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$ is a weight $B$-module with weight space $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu = (M_\mu)^*$

Given a map $\varphi : M_\mu \to \mathbb{C}[z]$, $(h - \mu z)\varphi(m) = \varphi((h - \mu z)m) = 0$, hence $(M_\mu)^* \subset (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu$.Therefore: $\sum\limits_{\lambda\in\mathbb{C}} (M_\mu)^* \subset \sum\limits_{\lambda\in\mathbb{C}} (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu \subset (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$.

By definition $\sum\limits_{\lambda\in\mathbb{C}} (M_\mu)^* = (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$. Therefore: $\sum\limits_{\lambda\in\mathbb{C}} (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu = (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$.

Each $M_\lambda \neq 0$ is free of finite rank, hence $(M_\lambda)^*$ is free of finite rank and $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$is a free $\mathbb{C}[z]$-module, hence $z$-torsion free.

It follows $\sum\limits_{\lambda\in\mathbb{C}} (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu = \underset{\lambda\in\mathbb{C}}{\oplus} (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu$.

We have proved $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$is a weight space with $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^{**} \cong \underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda$.

We have proved the following:

**Theorem 11.** *Let $\mathcal{W}_B$ be the full subcategory of $\mathcal{O}_B$ consisting of the $z$-torsion free weight $B$-modules. Then there is a duality $(-)^* : \mathcal{W}_B \to \mathcal{W}_B$ given by: $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^* = \underset{\lambda\in\mathbb{C}}{\oplus} (M_\lambda)^*$, where $(\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu = (M_\mu)^*$ and $\underset{\lambda\in\mathbb{C}}{\oplus} (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*_\mu = (\underset{\lambda\in\mathbb{C}}{\oplus} M_\lambda)^*$.*

# References

[An] Anick D.J. On the homology of associative algebras, Transactions of the American Mathemtaical Society, Vol. 296, No. 2, pp. 641-659, (1986).

[AR] Auslander, M.; Reiten , I.; Representation theory of artin algebras IV, Invariants given by almost split sequences, Communications in Algebra, 5(5), 443-518 (1977).

[BGG] Bernstein I., Gelfand, I. Gelfand, S. A certain category of $\mathfrak{g}$-modules, Funckcional. Anal. i Prilozen. **10**, 2, pp. 1-8, (1976).

[Co] Coutinho S.C. A Primer of Algebraic D-modules, London Mathematical Society, Students Texts 33, 1995

[Da] Dade, E. C. Group-graded rings and modules. Math. Z. 174 (1980), no. 3, 241–262.

[EW] Erdmann K. Wildon M. J. Introduction to Lie Algebras, Springer Undergraduate Mathematics Series, 2006.

[GM1] Green, E. L.; Martínez Villa, R.; Koszul and Yoneda algebras. Representation theory of algebras (Cocoyoc, 1994), 247–297, CMS Conf. Proc., 18, Amer. Math. Soc., Providence, RI, 1996.

[GM2] Green, E. L.; Martínez-Villa, R.; Koszul and Yoneda algebras. II. Algebras and modules, II (Geiranger, 1996), 227–244, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.

[LbS]Le Bruyn, Lieven; Smith, S. P. Homogenized sl(2).Proc. Amer. Math. Soc. 118 (1993), no. 3.

[MZ] Martínez-Villa, Roberto; Zacharia, Dan; Approximations with modules having linear resolutions. J. Algebra 266 (2003), no. 2, 671–697.

[MB] MacLane S. Birkhoff G. Algebra, AMS Chelsea Publishing, 1999.

[Ma] Mazorchuk, V. Lectures on $s\ell_2(\mathbb{C})$-modules. Imperial College Press, World Scientific Publishing Co. Pte. Ltd. 2010.

[MMo1] Martinez-Villa R. Mondragon J. On the homogenized Weyl algebra, Preprint, Centro de Ciencias Matematicas UNAM, (2012).

[MMo2] Martinez-Villa R. Mondragon J. Skew group algebras, invariants and Weyl Algebras, Preprint, Centro de Ciencias Matematicas UNAM, (2012).

[Mil] Miličić, D. Lectures on Algebraic Theory of D-modules. University of Utah 1986.

[P] Popescu N.; Abelian Categories with Applications to Rings and Modules;, L.M.S. Monographs 3, Academic Press 1973.

[Sm] Smith, P.S.; Some finite dimensional algebras related to elliptic curves, Rep. Theory of Algebras and Related Topics, CMS Conference Proceedings, Vol. 19, 315-348, Amer. Math. Soc 1996.

[Y] Yamagata, K.; Frobenius algebras. Handbook of algebra, Vol. 1, 841–887, North-Holland, Amsterdam, 1996.

Centro de Ciencias Matematicas, UNAM Morelia.
*E-mail address*: `mvilla@matmor.unam.mx`
*URL*: `http://www.matmor.unam.mx`